

Avifauna of the Pilón Lajas Biosphere Reserve and Communal Lands, Bolivia

A. BENNETT HENNESSEY, SEBASTIAN A. HERZOG, MICHAEL KESSLER
and DANIEL ROBISON

Summary

Pilón Lajas Biosphere Reserve and Communal Lands (Pilón Lajas) ranges from 300 m to 1,400 m altitude in the departments of Beni and La Paz, Bolivia. It is located in the Bolivian and Peruvian lower yungas Endemic Bird Area (054) and within one of the regions of highest conservation priority in the Neotropics. Pilón Lajas includes primary evergreen tropical lowland forest, foothill forest, and lower montane forest. Vegetation heterogeneity forms the basis for the high species richness of the area. A population of approximately 600 Chiman, Moseten and Tacana indigenous people live within the reserve. Little has been published regarding the avifauna of Pilón Lajas. Here we present the results of visual observations and tape recordings of 22 ornithological surveys, recording 502 bird species. Serranía Pilón is the most intensively studied and species-rich mountain range with 332 species, 48 of which have been recorded only in this area within Pilón Lajas. We present 26 new departmental records and evidence of breeding for 42 species. The reserve protects three globally Vulnerable and three globally Near Threatened species, four species of High Conservation Priority and 32 of Medium Conservation Priority. Together with Madidi National Park and the adjacent Bahuaja Sonene National Park in south-eastern Peru, Pilón Lajas forms a unique protected area system of more than 3.5 million ha and plays an important role in the establishment of biological corridors along the eastern Andean slope in Bolivia.

Introduction

Pilón Lajas Biosphere Reserve and Communal Lands (*Reserva de la Biósfera – Tierra Comunitaria de Origen Pilón Lajas* in Spanish, hereafter referred to as Pilón Lajas) covers c. 400,000 ha on the Andean foothills in north-west Bolivia (Figure 1). It is located in the Bolivian and Peruvian lower yungas Endemic Bird Area (EBA 054) of Stattersfield *et al.* (1998) and within one of the regions of highest conservation priority in the Neotropics (Stotz *et al.* 1996). Initially proposed in 1974, Pilón Lajas was recognized as a Biosphere Reserve by UNESCO in 1977 and as a Biosphere Reserve and Indigenous Territory by the Bolivian Government in 1992 (Sherholtz 1997). The area was legally designated but had no physical protection until 1994, when Veterinarians Without Frontiers (VSF) received funding from the European Union to design an operational plan for the reserve. Through an agreement with the Bolivian National Protected Areas Service, VSF has been co-administering Pilón Lajas since 1996.

Pilón Lajas is in an area with a moderate level of historic and current human activity and settlement. A population of approximately 600 Chiman, Moseten




Figure 1. Map showing Pilón Lajas Biosphere Reserve, Bolivia.

and Tacana indigenous people live within the reserve. The Beni River, the western border of Pilón Lajas (Figure 1), was the principal "highway" between the Andes and the rubber/brazil nut forests to the north for a century until a new road was completed in 1989. This new road, forming Pilón Lajas' eastern border and connecting the area to the city of La Paz, was used for a government-sponsored colonization programme of families from other parts of the country. Many of these colonists believe they have a right to land within Pilón Lajas. Although parts of the reserve are extremely remote, requiring expeditions of several days to be reached, other parts have long been accessible. It is estimated that only 1% of Pilón Lajas has been deforested for agriculture in the last century, but perhaps 80% of the lower tropical forest has been heavily hunted during the alligator hide and wildlife skin boom in the 1960s and 1970s. Between 1980 and 1996, the mahogany lumber boom heavily affected the same area, particularly within 10 km of rivers and large streams. Up until 1996, five lumber companies with dubious legal status and approximately 300 chain sawyers selectively logged and hunted in the reserve.

The Pilón Lajas management plan has subdivided the reserve into specific zones to combine indigenous community rights with conservation initiatives. Since 1996, a core of 20 park guards (many belonging to local indigenous groups) has been hired and trained and four park guard stations have been established in highly sensitive areas along borders to protect the reserve.

Table 1. Survey areas visited by 22 ornithological expeditions in Pílon Lajas, Bolivia, between 1989 and 1999.

| Survey area | Elevational range (m) | Number of visits |
|----------------------|-----------------------|------------------|
| Lowland forest areas | 350–500 | 5 |
| Serranía Bala | 450–800 | 2 |
| Serranía Beu | 500–1,400 | 4 |
| Serranía Cuchilla | 450–1,000 | 2 |
| Serranía Pílon | 400–1,100 | 8 |
| Serranía Zaguacales | 400–800 | 1 |

The aim of this paper is to bring together unpublished avifaunal information into an extensive document that can be used for informed conservation decision making and land protection priority assessments. This is important for the protection and global attention of Pílon Lajas and the general Andean protected area corridor of Madidi and Bahuaia Sonene National Parks.

Methods

This paper is based on 22 unpublished ornithological surveys by 15 researchers over 148 days between 1989 and 1999 (Table 1, Appendix 1) in Pílon Lajas. The principal survey methods were non-systematic visual and acoustic observations as well as tape recordings of vocalisations. Limited mist-netting was carried out by the first author (ABH) in Serranía Cuchilla in 1999 and by F. Osorio and others (Colección Boliviana de Fauna, La Paz; Appendix 1). A small number of specimens were collected by F. Osorio and others and deposited at the Colección Boliviana de Fauna.

Altitude ranges and abundance estimates are based on survey results by ABH. ABH surveyed different points at least 200 m apart, arriving before sunrise and identifying and sound-recording vocalizations of the dawn chorus. Thereafter, different trails were surveyed, covering 1–3 km, often pausing work between 12h00 and 15h00. ABH observed, sound-recorded and verified identifications with the use of prerecorded tapes or playback, including rebound playback (broadcasting the first response to playback). Each evening ABH completed a checklist of the day's observations, noting estimated abundances, field time, distance walked and evidence (sight records, heard or tape-recorded). Abundance estimates were classified as; common, recorded (visual observation or auditory) daily in small numbers (> 5); fairly common, recorded daily in small numbers (< 5); uncommon, recorded once in three days; and rare, recorded once during the survey. Tape-recording bird vocalizations is the most effective method for rapidly surveying neotropical forest avifaunas (Parker 1991; see also Haselmayer and Quinn 2000) and is an accurate means of species identification that can be easily referenced. Over 400 recordings by ABH from Pílon Lajas have been archived at the Cornell Laboratory of Ornithology, Macaulay Library of Natural Sounds (MLNS). Fieldwork covered all months except for January, March and May, and included all habitat types within Pílon Lajas and as much area as possible with the largest gap between sites being 15 km. For the purpose of vegetation classification, we used the elevational belts of Stotz *et al.* (1996): lower

tropical forest (< 500 m), hill tropical forest (500–900 m) and upper tropical forest (900–1,600 m). This was modified as necessary by reference to topographical location (level areas, slopes, ridges, etc.), geological substrate and state of vegetation succession. It should be noted that Parker (1989) erroneously gave 1,200 m as the highest altitude of the road crossing Serranía Pilón south of Yucumo (Figure 1). Rather, as indicated by Parker *et al.* (1991) and as confirmed by our measurements, the highest point of this road is located at *c.* 1,000 m.

Study Area

Pilón Lajas is situated at 14°26' to 15°28'S and 68°42' to 68°02'W in Beni and La Paz departments with an altitudinal range from 300 m to 1,400 m, although there probably are a few peaks up to 1,600 m (detailed topographical maps of the area are unavailable). These mountain ranges are relatively isolated from the main Andean ridges. The nearest range of similar altitude is about 35 km to the south-west and the closest peaks above 1,400 m are about 75 km to the south of Pilón Lajas.

The vegetation of Pilón Lajas is quite varied, changing in relation to humidity, elevation and geological substrate. At Rurrenabaque (14°26'S 68°31'W, 227 m), just north of Pilón Lajas (Figure 1), mean annual precipitation is 2,150 mm with a marked peak from November to April and a mean annual temperature of 25.8 °C. The natural vegetation in about 38% of the reserve corresponds to evergreen tropical lowland forest. Higher elevations in Pilón Lajas, particularly those exposed to incoming clouds from the north-east, are somewhat cooler, receive considerably more precipitation and, above *c.* 1,000 m, are frequently shrouded in mist. Here, forests are covered in epiphytic mosses, liverworts, ferns, bromeliads and orchids, and trees are often stunted.

Geologically, the area consists mainly of schists, loams and white sandstones of Tertiary age. Flat valley bottoms with mixed alluvial, nutrient-rich soils and good water supply are periodically flooded in the rainy season and support 30–40 m tall forest with abundant palms (e.g. *Iriartea deltoidea*, *Socratea exorrhiza*), figs and leguminaceous trees. Gentle to fairly steep slopes are covered with forest of somewhat lower stature with up to 150 tree species and about 200 species of vascular epiphytes per ha (Seidel 1995, Smith and Killeen 1998, T. Krömer pers. com.). Above about 1,000 m the tree flora is dominated by the families Lauraceae, Leguminosae and Melastomataceae. Locally, there are dense stands of mahogany *Swietenia macrophylla*, many of which have been exploited in the last decades. Areas with white sand soils are very poor in nutrients and support a distinct and quite species-rich flora (Kessler 2001). Steep slopes, particularly at higher elevations, are characterized by unstable soils and frequent landslides and thus often support vegetation in different stages of succession, ranging from low, dense thickets to young, 10–15 m tall forest with many slender trees. The highest ridges often have shallow, leached soils and stunted, species-poor forest.

The overall picture emerging from the above combination of ecological factors is a striking heterogeneity of the vegetation that is conspicuous in the field (particularly from the air) and that forms the basis for the high biological species richness of the area. Botanically, individual vegetation types are of about average diversity compared with other Neotropical areas, but the high number of distinct vegetation types leads to an exceedingly diverse overall flora that contains

numerous elements not known elsewhere in the Bolivian Andes (Smith and Killeen 1998, Kessler 2001).

Results

Currently, 502 bird species are known from Pilón Lajas (Appendix 2), more than one third of the 1,398 species reported in Bolivia (Hennessey *et al.* 2003). We believe that this includes at least 90% of the resident bird species in the reserve. Whereas 309 species (62%) were recorded in the approximately 150,000 ha of lower tropical forest in Pilón Lajas, only 79 species were exclusively found in this zone. The lower tropical forest of Pilón Lajas appears quite species poor in comparison with other nearby lowland localities such as Alto Madidi, departamento (dpto.) La Paz, with 405 species (Parker and Bailey 1991) and the 5,000 ha Tambopata-Candamo Reserved Zone, Peru, with 572 species (Parker *et al.* 1994). This relatively low diversity may well be a result of the geographical position of Pilón Lajas on the southern edge of the Amazon basin where tropical lowland forest is limited to a narrow, 10–30 km wide belt along the Andean foothills.

Serranía Pilón is the most intensively studied and species-rich range within Pilón Lajas with 332 species, 48 of which have only been recorded in this area. Abundances of birds appear to be higher on Serranía Pilón than on comparable nearby ranges. Daily mixed-species flock encounters on serranías Bala and Cuchilla (Figure 1) as well as on Serranía Sadiri in Bolivia's Madidi National Park (ABH pers. obs.) ranged between one and three per day, whereas encounter rates on Serranía Pilón were between three and five per day. This is probably due to a higher level of precipitation in the area and the unusual topography of Serranía Pilón with several peaks, opposed to a typical single-peaked range, creating more habitat area between 600 m and 900 m.

Upper tropical forest exists in Pilón Lajas only on serranías Bue, Chepete and Muchanes. All surveys in this habitat were restricted to Serranía Beu, where 246 bird species have been recorded, 46 of which were only found in upper tropical forest. Serranía Beu is the only area in Pilón Lajas with large patches of bamboo forest between 1,200 to 1,400 m; they cover an area of approximately 300 ha.

With respect to Arribas *et al.* (1995) and Remsen and Traylor (1989), the Pilón Lajas bird list includes 26 new records for dpto. Beni and three for dpto. La Paz (Appendix 2). However, these new records by no means represent true range extensions for the respective species. Bolivian departments are formed by arbitrary political borders that have little or no value in a habitat or conservation context. For example, about 85% of the area of the department of Beni is covered by pampas (savannah grasslands) whereas less than 1% contains hill tropical forest, from where most of our new records originated.

Pilón Lajas is located approximately 25 km south of a large expanse of pampas, and its northern and eastern border is formed by a main road. Through the years, the area has seen the establishment of pampas bird species as colonists have cleared forest. In Rurrenabaque, each year more open-country species are observed as the 30 km wide forest corridor between the town and the true pampas is being settled and destroyed. With continued forest clearing activities an increasing number of open-country species and austral migrants can be

expected in open areas of Pilón Lajas with many of the former probably becoming resident along the edge of the reserve.

The majority of birds in Pilón Lajas increase their territorial singing in September/October. The full onset of the breeding season is linked with the beginning of the rainy season in November. During the rainy season there is a marked increase in insect abundance and more trees bear fruit. Local hunters claim that large mammals have more fat from February to April. We found reproductive evidence for 42 species (Appendix 2). Nest-building activity was observed between late July and mid-November (18 species), nesting activity (egg and hatchling stages) from mid-September to early November (nine species) and recent fledglings (15 species) between early October and mid-February (with two exceptions in mid-July).

Mixed-species flocks

ABH observed 45 mixed-species foraging flocks containing a total of 102 species in hill tropical forest (500–900 m). In the tropical lowlands, foraging flocks are often strictly separated into canopy and understorey flocks (e.g. Munn 1985, Jullien and Thiollay 1998). Due to the variable nature of hill tropical forest with steep slopes, landslides, cliffs, forest edges, stunted forest and tall forest, flock formations in Pilón Lajas were loosely categorized into three types: omnivorous tanager flocks, mid-canopy and understorey insectivorous mixed-species flocks.

The tanager canopy flocks were fast-moving, omnivorous flocks of the upper canopy based around *Tangara* species with Paradise Tanager *Tangara chilensis* as the nuclear species and Green-and-gold Tanager *T. schrankii*, Turquoise Tanager *T. mexicana* and Blue Dacnis *Dacnis cayana* as other characteristic species. Mid-canopy insectivorous flocks ranged from c. 5 m above ground to well inside the highest canopy. These flocks were slower moving and sometimes large (c. 40 individuals) with most species foraging for insects. Characteristic species were Slate-throated Redstart *Myioborus miniatus*, Black-spotted Barbet *Capito niger*, Masked Tityra *Tityra semifasciata*, Guira Tanager *Hemithraupis guira*, Bronze-green Euphonia *Euphonia mesochrysa*, Bay-headed Tanager *Tangara gyrola* and Tropical Parula *Parula pitiayumi*. The nuclear species of understorey insectivorous flocks was Red-crowned Ant Tanager *Habia rubica*, with Buff-throated Foliage Gleaner *Automolus ochrolaemus*, Carmiol's Tanager *Chlorothraupis carmioli*, Chestnut-tailed Antbird *Myrmeciza hemimelaena*, McConnell's Flycatcher *Mionectes macconnelli* and Tawny-crowned Greenlet *Hylophilus ochraceiceps* as other characteristic species.

However, the majority of species in these flocks in hill tropical forest were not exclusive or even particular to any of the three flock types. This strongly contrasts with the rigidly organized flocks of Amazonian lowlands (Munn 1985, Jullien and Thiollay 1998) and supports the notion that conclusions on mixed-species flock structure and organization drawn from Neotropical lowland forests cannot be generalized to all tropical forests (Jullien and Thiollay 1998, Herzog *et al.* submitted MS).

Species Accounts

Orinoco Goose *Neochen jubata*

This Near Threatened species has been observed only once in Pilón Lajas at an

unknown date in 1994 by F. Osorio and others, although indigenous people say the bird is sporadically seen on Rio Quiquebey. If a population still exists in Pilón Lajas, it will be in isolated regions away from hunters.

Harpy Eagle *Harpia harpyja*

This globally Near Threatened species has never been sighted in Pilón Lajas by trained observers. However, the Chiman and Mosesten indigenous people are familiar with the bird and call it Bonij, but they do not have a name for the similar Crested Eagle *Morphnus guianensis*. ABH was shown a photo of a shot Harpy Eagle taken by an unknown tourist on the shore of the Beni River within Pilón Lajas around 1970, proving that it did exist in the area, but the exact location of the dead bird was unclear. A Mosesten elder reported that a family of Harpy Eagles exists east of S. Beu and described the species accurately as he said they ate one bird five years ago while hunting for White-bellied Spider Monkeys *Ateles belzebuth*. A Chiman hunter reported the location of a nest in Pilón Lajas south-west of the mouth of the Suapi River. We believe that a handful of Harpy Eagles still exists within Pilón Lajas, but that this population is threatened and restricted to areas with low human impact.

Southern Horned Curassow *Pauxi unicornis*

Parker (1989) gave the following account of this globally Vulnerable species from a 1989 survey in Serranía Pilón along the Caranavi-Yucumo road at the south-eastern border of Pilón Lajas: "A man who has lived for one year near the road pass on the Serranía del Pilón accurately described this species and was certain that he had seen (eaten?) it in ridgetop forest several times. Called "mutum"." This record would appear to be a communication error as the name mutún locally refers to Razor-billed Curassow *Mitu tuberosa*.

None of the 22 expeditions in Pilón Lajas, including two to inaccessible, pristine and apparently suitable habitat for this species between serranías Beu and Chepete, has resulted in a single observation of this species. Likewise, numerous interviews including a survey of local indigenous knowledge by ABH and A. Perry have failed to produce any indication that the species has ever occurred in the reserve. We believe that *Pauxi unicornis* does not exist in Pilón Lajas.

Oilbird *Steatornis caripensis*

Oilbird is a gregarious species, breeding colonially in caves. A single cave can hold up to 1,000 individuals (Thomas 1999; Duguid *in litt.* 2000). Only eight cave sites are known in Bolivia (ABH pers. obs.). Oilbirds could be threatened by habitat loss and scarcity of isolated breeding areas. On 4 November 1998, ABH and park guards observed 350 individuals leaving an inaccessible cave at 600 m on the edge of Serranía Chepete, beside the Beni River, during sunset and found fresh eggshells below the entrance to the cave.

Lanceolated Monklet *Micromonacha lanceolata*

One immature and two adults were observed for approximately 15 min on 9 December 1995 by ABH and A. Perry in the subcanopy 8 m above ground at

1,150 m on Serranía Beu, dpto. La Paz. One adult fed a small unidentified insect to the immature and then perched 30 cm away while the second adult landed beside the immature and gave it a katydid-like insect longer than its bill. The immature clasped the insect in its bill for approximately 2 min before swallowing it. The immature's breast had less clearly defined black streaks and greyer mottling than both adults. These birds were observed on a steep forested slope with many successional stages from frequent landslides to dense forest patches with young trees. Another individual was observed in Pílon Lajas by A. Perry and R. Mitchell (pers. comm.) on the eastern edge of Serranía Beu at 1,000 m on 3 December 1996. These are the second and third observation of Lanceolated Monklet in Bolivia (Arribas *et al.* 1995). The first record was obtained by S. Cardiff and J.V. Remsen (*in litt.* 1998) at Cerro Asunta Pata (15°03'S 68°28'W), dpto. La Paz, in 1993. The species was also observed in Carrasco National Park, dpto. Cochabamba, in July 1998 by A. Mee (*in litt.* 1998).

Scaled Fruiteater *Ampelioides tschudii*

Two birds, possibly a pair were noted repeatedly on Serranía Cuchilla, dpto. Beni, at 900 m in 1998 and 1999, especially by voice (MLNS 101799, 101819, 101874, 101877; Mayer 2000). On 17 November 1999 a male was observed by ABH and M. Olivera feeding an immature an orange fruit of approximately 2.5 cm in diameter and regurgitated half bill-sized black fruits. Previously known in Bolivia only from Serranía Bellavista (38 km N Caranavi), dpto. La Paz (Parker *et al.* 1980), and Río Tuichi Valley, dpto. La Paz (Perry *et al.* 1997). ABH also recorded calls of one individual (MLNS 101845) at 900 m on Serranía Sadiri (14°13'S 67°56'W) in Madidi National Park on 30 November 1999.

Subtropical Pygmy Owl *Glacidium parkeri*

C. König identified the first record of this species for Bolivia through recordings made by T.A. Parker during a 1979 expedition by the Museum of Natural Science, Louisiana State University, Baton Rouge, to Serranía Bellavista, dpto. La Paz (T.S. Schulenburg pers. comm.). B.M. Whitney (*in litt.* 2001) obtained the second record for Bolivia on Serranía Eslabon in Madidi National Park in August 1995 and the third in Pílon Lajas on Serranía Pílon at 700 m on 9 August 1997. Subsequently the species was recorded by ABH (MLNS 101656, 101689) several times on both the western (1,300 m) and eastern (1,200 m) slope of Serranía Beu and once at 750 m on Serranía Tequeje, Madidi National Park, on 6 April 2000.

Bolivian Recurvebill *Simoxenops striatus*

Until recently, this globally Vulnerable species was known from only three Bolivian localities (Parker *et al.* 1992), and Parker *et al.* (1996) considered it a high conservation and research priority. ABH observed and recorded several individuals on Serranía Beu at 800 m to 1,250 m from 17 to 24 September 1998 (MLNS 101637) and on Serranía Cuchilla at 800 m between 16 and 19 October 1998 (MLNS 101778; Mayer 2000), which represent two new localities for the species in Bolivia. Additionally, it has been found at three other new sites in dptos. La

Paz, Cochabamba and Santa Cruz by SKH and MK in 1996 and 1997. Further details will be published elsewhere.

Yungas Antwren *Myrmotherula grisea*

A single female-plumaged bird of this globally Vulnerable Bolivian endemic was seen by SKH in an understorey mixed-species flock at c. 850 m on Serranía Pílon on 4 August 1997, which represents the only record for Pílon Lajas. The bird was characterized by unmarked wing-coverts and warm olivish-brown upperparts with a slight rufous tinge on wings and tail (in contrast to the all-grey upperparts of the female Grey Antwren *M. menetriesii*) and it resembled a female Yungas Antwren seen and tape-recorded by SKH together with a male (both singing) in Amboró National Park (dpto. Santa Cruz). Despite the presence of apparently suitable habitat within the species's elevational range, Yungas Antwren appears to be rare in Pílon Lajas.

Phyllomyias sp.

This undescribed species was first observed and tape-recorded in June 1989 at 850–900 m on Serranía Pílon along the Caranavi–Yucumo road (Parker *et al.* 1991), who tentatively identified the birds as Planalto Tyrannulets *Phyllomyias fasciatus*. However, they noted that their vocalizations were faster and higher pitched than those of birds in south-east Brazil and north-east Argentina and concluded that “the Serranía Pílon population may represent an undescribed form”. On 2 June 1997 SKH tape-recorded three unidentified *Phyllomyias* tyrannulets in the canopy of humid forest edge at 1,150 m at Cerro Asunta Pata, dpto. La Paz (15°03'S 68°28'W), and this recording was later identified with the help of B.M. Whitney as belonging to the same undescribed *Phyllomyias* taxon as recorded by Parker and Gell-Mann on Serranía Pílon (see Mayer 2000). On 3 September 1998, SKH and J.A. Balderrama collected two specimens at Cerro Asunta Pata and rediscovered the Serranía Pílon population at 850–1,000 m along the Caranavi–Yucumo road on 8 October 1998. Several tape-recordings from both localities show that the vocal differences noted by Parker *et al.* (1991) between Andean and south-east Brazilian populations are both consistent and significant (recordings of *P. fasciatus* from south-east Brazil provided by J. Mazar Barnett). A formal description of the species is currently in preparation (Herzog unpublished data).

It is noteworthy that despite the presence of apparently suitable hill tropical forest habitat on Serranías Beu, Cuchilla and Pílon the species has so far only been found at a single locality in Pílon Lajas, indicating that it may have a naturally patchy distribution and a small range.

Shrike-like Cotinga *Laniisoma elegans*

This globally Vulnerable species was observed by ABH on 22 September 1999 15 m above the ground on the northern side of Serranía Beu in steep-sloped hill tropical forest (850 m). This is only the second observation for Bolivia of this poorly known species (Hennessey *et al.* 2003).

Scarlet Tanager *Piranga olivacea*

This Nearctic-Neotropical migrant has suffered population declines attributed to habitat loss in its wintering range (McNair and Escobar 1993). However, its non-breeding distribution in South America is poorly known (Parker *et al.* 1996). ABH observed this species on Serranía Pílon in November 1996 and February 1997 and 1998, on Serranía Cuchilla in November 1999, and on Serranía Sadiri in Madidi National Park in November 1999. Wintering males were identified by their contrasting black wings in November, and by the splotchy breeding plumage in February. The identification of females was assumed through their presence with Scarlet Tanager males, which in February consisted of associating groups of 5–8 individuals. The species was fairly common in all years and areas. For example, on a total of seven days on Serranía Pílon during the months stated above, ABH daily observed between two and seven individuals in non-breeding plumage in mid-canopy foraging flocks of hill tropical forest (600–900 m). We believe these records from serranías Pílon, Cuchilla and Sadiri indicate that a considerable portion of the species' population winters along the humid foothills of the Bolivian Andes. This habitat is still largely unaltered in Bolivia; hence, a decline due to habitat loss is probably largely caused by the alteration of habitat in the breeding range or along the migration route of Scarlet Tanager.

Buff-rumped Warber *Basileuterus fulvicauda*

J. Ohlson (pers comm.) observed several individuals on Serranía Bala from 25 to 28 April 1999 and recorded one (Mayer 2000). Observations and recordings (MLNS 87814 and 101816) by ABH and SKH on serranías Pílon and Cuchilla have exclusively been of the similar River Warbler *Basileuterus rivularis*. The song of both species is difficult to differentiate. Thus, both species occur within at most 30 km of each other in Pílon Lajas, but the actual contact zone as well as the degree of sympatry and hybridization (if any) remain to be determined.

Discussion

Recorded from Pílon Lajas are four species of Bolivian endemics, eight species of range-restricted endemics (Stattersfield *et al.* 1998), 28 zoogeographical regional endemics (Stotz *et al.* 1996), three Vulnerable and three Near Threatened species (BirdLife International 2000), and four High Conservation Priority and 32 Medium Conservation Priority species (Parker *et al.* 1996) (Appendix 2, Table 2). Of the High and Medium Conservation Priority species, eight have their centre of abundance in lower tropical (< 500 m), nine in hill tropical (500–900 m), 17 in upper tropical (900–1,600 m) and two in middle tropical (1,600–2,600 m) forest. All High Conservation Priority species in Pílon Lajas appear to be rare, with only one record per species, except for *Simoxenops striatus*. These apparently extremely low population densities imply that optimal conditions for these species may not be met in Pílon Lajas despite the presence of apparently suitable habitat and that the reserve alone cannot warrant their effective conservation in Bolivia.

Pílon Lajas protects a healthy part of the Bolivian and Peruvian lower Yungas EBA (054) (Stattersfield *et al.* 1998). Eight range-restricted species have been

Table 2. Species in Pílon Lajas, Bolivia, listed as High and Medium Conservation Priorities by Parker *et al.* (1996) and Vulnerable (*) or Near Threatened (**) by BirdLife International (2000). Research priority and centre of abundance after Parker *et al.* (1996). Numbers in parentheses refer to the number of records.

| Species | Conservation priority | Research priority | Centre of abundance | Pílon Lajas abundance |
|-----------------------------------|-----------------------|-------------------|---------------------|-----------------------|
| Orinoco Goose** | | | | |
| <i>Neochen jubata</i> | High | High | Lower tropical | Very rare (1) |
| Fasciated Tiger Heron | Medium | High | Hill tropical | Very rare (2) |
| <i>Tigrisoma fasciatum</i> | | | | |
| Solitary Eagle | Medium | Medium | Upper tropical | Very rare (2) |
| <i>Harpyhaliaetus solitarius</i> | | | | |
| Harpy Eagle** | Medium | High | Lower tropical | Very rare (?) |
| <i>Harpia harpyja</i> | | | | |
| Red-throated Caracara | Medium | Medium | Lower tropical | Fairly common |
| <i>Ibycter americanus</i> | | | | |
| Maroon-chested Ground Dove | Medium | Medium | Middle montane | Very rare (1) |
| <i>Claravis mondetoura</i> | | | | |
| Blue-and-yellow Macaw | Medium | Medium | Lower tropical | Fairly common |
| <i>Ara ararauna</i> | | | | |
| Military Macaw* | Medium | Medium | Upper tropical | Uncommon |
| <i>Ara militaris</i> | | | | |
| Red-billed Parrot | Medium | Medium | Upper tropical | Uncommon |
| <i>Pionus sordidus</i> | | | | |
| Rufous-vented Ground Cuckoo | Medium | Medium | Lower tropical | Very rare (1) |
| <i>Neomorphus geoffroyi</i> | | | | |
| Band-bellied Owl | Medium | Medium | Hill tropical | Fairly common |
| <i>Pulsatrix melanota</i> | | | | |
| Silky-tailed Nightjar | Medium | Medium | Lower tropical | Very rare (1) |
| <i>Caprimulgus sericocaudatus</i> | | | | |
| White-browed Hermit | Medium | Medium | Hill tropical | Very rare (1) |
| <i>Phaethornis stuarti</i> | | | | |
| Rufous-crested Coquette | Medium | Medium | Upper tropical | Very rare (1) |
| <i>Lophornis delattrei</i> | | | | |
| Many-spotted Hummingbird | Medium | Medium | Hill tropical | Very rare (1) |
| <i>Taphrospilus hypostictus</i> | | | | |
| Crested Quetzal | Medium | Medium | Upper tropical | Fairly common |
| <i>Pharomachrus antisianus</i> | | | | |
| Ash-browed Spinetail | Medium | Medium | Upper tropical | Rare |
| <i>Cranioleuca curtata</i> | | | | |
| Bolivian Recurvebill* | High | High | Hill tropical | Uncommon |
| <i>Simoxenops striatus</i> | | | | |
| Yungas Antwren* | High | High | Hill tropical | Very rare (1) |
| <i>Myrmotherula grisea</i> | | | | |
| Rough-legged Tyrannulet | Medium | Medium | Hill tropical | Uncommon |
| <i>Phyllomyias burmeisteri</i> | | | | |
| Spectacled Bristle Tyrant | Medium | Medium | Hill tropical | Uncommon |
| <i>Phylloscartes orbitalis</i> | | | | |
| White-bellied Pygmy Tyrant | Medium | Medium | Hill tropical | Fairly common |
| <i>Myiornis albiventris</i> | | | | |
| Yungas Tody Tyrant | Medium | Medium | Upper tropical | Common |
| <i>Hemitriccus spodiops</i> | | | | |
| Buff-throated Tody Tyrant** | Medium | Medium | Upper tropical | Rare |
| <i>Hemitriccus rufifigularis</i> | | | | |
| Shrike-like Cotinga | High | High | Upper tropical | Very rare (1) |
| <i>Laniisoma elegans</i> | | | | |

Table 2. Continued.

| Species | Conservation priority | Research priority | Centre of abundance | Pilón Lajas abundance |
|---|-----------------------|-------------------|---------------------|-----------------------|
| Scaled Fruiteater <i>Ampelioides tschudii</i> | Medium | Medium | Upper tropical | Uncommon |
| Andean Cock-of-the-Rock <i>Rupicola peruviana</i> | Medium | Medium | Upper tropical | Fairly common |
| Orange-eared Tanager <i>Chlorochrysa calliparaea</i> | Medium | Medium | Upper tropical | Very rare (1) |
| Pale-winged Trumpeter <i>Psophia leucoptera</i> | Medium | Low | Lower tropical | Rare |
| Red-and-green Macaw <i>Ara chloroptera</i> | Medium | Low | Lower tropical | Fairly common |
| Upland Antshrike <i>Thamnophilus aroyae</i> | Medium | Low | Upper tropical | Fairly common |
| Scaled Antpitta <i>Grallaria guatemalensis</i> | Medium | Low | Upper tropical | Rare |
| Slaty Gnateater <i>Conopophaga ardesiaca</i> | Medium | Low | Upper tropical | Uncommon |
| Slaty Tanager <i>Creurgops dentata</i> | Medium | Low | Middle montane | Very rare (1) |
| Bronze-green Euphonia <i>Euphonia mesochrysa</i> | Medium | Low | Upper tropical | Common |
| Golden-eared Tanager <i>Tangara chrysotis</i> | Medium | Medium | Upper tropical | Very rare (2) |

recorded of a potential 13 species found in Bolivia. The range-restricted endemism of Serranía Beu demonstrated the area to be a potential Important Bird Area. Serranía Beu and the surrounding area are sufficiently difficult to access through climbing, long hikes and the lack of road access, that the areas may be easily protected from habitat destruction through local population pressure.

Serranía Pilón is an important area in Pilón Lajas to protect foothill-specialized species. The foothill forest, though not considered specialization enough to deserve EBA status, is a unique ecosystem that deserves conservation attention. Through high levels of precipitation and the mix between lowland and Yungas species, these areas tend to be the richest sites of avian diversity on the planet. With no avifaunal data from Serranía del Tigre (13°38'S 68°40'W), mostly outside the Madidi National Park border, Serranía Pilón is the richest area of foothill forest in Bolivia. The site, with a main road travelling through the area, also contains ecotourism-sustainable development possibilities.

Pilón Lajas has common borders with Madidi National Park to the north-west and with the Moseten Indigenous Area, which applies sustainable forestry practices, to the south. However, the north- and south-eastern borders of Pilón Lajas are adjacent to major roads that have enabled human settlement on the park boundaries, posing a serious future threat to the reserve. Pilón Lajas is trying to counteract this encroachment with two park guard stations along both roads and the implementation of public education programmes. Recently a sustainable ecotourism project has been designed with a local indigenous group at the mouth of Rio Quiquebey.

Together with Madidi National Park and the directly adjacent Bahuaaja Sonene

National Park in south-eastern Peru, Pilón Lajas forms part of a unique protected area system of more than 3.5 million ha. It further represents a vital link towards foothill and montane forest areas further south and plays an important role in the establishment of biological corridors along the eastern Andean slope in Bolivia.

Acknowledgements

We would like to thank Veterinarians Without Frontiers and the British, Dutch and Swiss Governments for their support and efforts to protect this special area of the earth and to support our insistence for more field investigations. ABH would like to especially thank Sjoerd Mayer for his meticulous review of Pilón Lajas sound recordings, the use of his pre-release CD-ROM *Birds of Bolivia 2*, and the hundreds of email messages on “mystery” sounds, queries and bird sound discussions. We are grateful to Bret M. Whitney for his assistance with Pilón Lajas research and for his time and patience with his remarkable identifications of “mystery” sounds that no one else could identify. ABH is most appreciative for the donations of tapes and equipment by Greg Budney at Cornell’s Macaulay Library of Natural Sounds. In addition, we thank Doug Stotz, Tom Schulenberg and Jon Fjeldså for their informative answers to the many questions involving Pilón Lajas research. Financial support was provided to ABH by Veterinarians Without Frontiers as well as the British, Dutch and Swiss Governments, and to MK by the Deutsche Forschungsgemeinschaft.

Appendix 1. Ornithological surveys in Pilón Lajas, Bolivia.

| Researchers | Site | Date | Period (days) |
|---|---|----------------|---------------|
| T.A. Parker | S. Pilón (600–1,000 m) | June 1989 | 4 |
| F. Osorio, S. Barrera, J.F. Guerra, J. Sarmiento, L. Villalba | Quiquibey and Suapi rivers and Colorado (350–500 m) | June 1993 | 10 |
| ABH | Suapi River (450 m) | November 1995 | 4 |
| A. Perry, ABH | S. Beu (western end, 700–1,300 m) | December 1995 | 7 |
| F. Sagot | Colorado (350–400 m) | June 1996 | 6 |
| ABH, R. Silva | Colorado to Quiquibey (350–400 m) | July 1996 | 7 |
| ABH | S. Bala (400–700 m) | October 1996 | 3 |
| ABH, R. Mitchell | S. Bala (500–900 m) | November 1996 | 4 |
| ABH | S. Pilón (600–1,000 m) | November 1996 | 7 |
| A. Perry, R. Mitchell | S. Zaguacales (350–700 m) | November 1996 | 6 |
| A. Perry, R. Mitchell | S. Beu (eastern end, 600–1,400 m) | December 1996 | 11 |
| ABH | S. Pilón (700–750 m) | February 1997 | 1 |
| ABH | S. Pilón (500–1,000 m) | April 1997 | 5 |
| ABH | S. Pilón (600–1,000 m) | June 1997 | 7 |
| SKH, MK | S. Pilón (450–1,000 m) | July 1997 | 18 |
| B.M. Whitney, ABH | S. Pilón (400–750 m) | August 1997 | 3 |
| ABH | S. Pilón (350–1,000 m) | February 1998 | 11 |
| ABH | S. Beu (western end, 1,200–1,450 m) | September 1998 | 7 |
| ABH | S. Beu (western end, 800–1,000 m) | September 1998 | 5 |
| SKH, J.A. Balderrama | S. Pilón (450–500 m) | October 1998 | 4 |
| ABH | S. Cuchilla (500–900 m) | October 1998 | 6 |
| ABH | S. Cuchilla (700–900 m) | November 1999 | 12 |

Appendix 2: List of Bird species from Pilon Lajas Biosphere Reserve and Communal Lands, Departments of Beni and La Paz, Bolivia, including evidence, sites, minimum and maximum elevational records and abundance estimates.

Nomenclature and taxonomic arrangement accord with Rensen and Taylor (1989) and Arribas et al. (1995).

Species: * Zoogeographical endemic (Stotz et al. 1996); **, Range restricted endemic (Stattersfield et al. 1998); ***, Political endemic.

Evidence (in order of priority): SP, specimen (Coleccion Boliviana de Fauna); T, sound recording (Cornell Library of Natural Sounds); O, visual observation; A, auditory; L, local information.

Sites: 1, tropical forest; 2, Serranía Zaguacales; 3, S. Bala; 4, S. Cuchilla; 5, S. Pilón; 6 S. Beu.

Min: Minimum elevation, L, Lowland (350–500 m).

Max: Maximum elevation, The highest viable forest habitat in Pilon Lajas is 1400 m.

Abundance: C, common, recorded (visual observation or auditory) daily in small numbers (> 5); F, fairly common, recorded daily in small numbers (< 5); U, uncommon, recorded once in 3 days; R, rare, recorded once in 6 days or less, VR, recorded 4 or less times within Pilon Lajas, number of records in quotes; Mn, Northern migrant; Ms, Southern migrant; Mm; migrant from the Andes; +, including resident population.

Breeding information: NA, attending nest; NB, building nest; NE, nest with eggs; NM, collecting nest material; F, fledgling.

| Species | Evidence | Sites | Min | Max | Abundance | Breeding Information |
|--|----------|---------|-----|-------|-----------|----------------------|
| <i>Tinamus tao</i> Grey Tinamou | T | 1–6 | L | 1,000 | C | |
| <i>Tinamus major</i> Great Tinamou | T | 1–6 | L | 600 | F | NE 17–10–98 |
| <i>Crypturellus cinereus</i> Cinereous Tinamou | A | 1,2,6 | L | 900 | R | |
| <i>Crypturellus soui</i> Little Tinamou | T | 1–6 | L | 900 | C | |
| <i>Crypturellus obsoletus</i> Brown Tinamou | T | 1,3–6 | L | 1,300 | C | |
| <i>Crypturellus undulatus</i> Undulated Tinamou | A | 1,2,4–6 | L | 800 | C | |
| <i>Tachybaptus dominicus</i> Least Grebe | O | 1 | L | 350 | VR (1Mn) | |
| <i>Phalacrocorax brasilianus</i> Neotropic Cormorant | O | 1,2,6 | L | 500 | F | |
| <i>Anhinga anhinga</i> Anhinga | O | 1 | L | 500 | F | |
| <i>Tigrisoma lineatum</i> Rufescent Tiger Heron | O | 1,3,5,6 | L | 700 | U | |
| <i>Tigrisoma fasciatum</i> Fasciated Tiger Heron | O | 1,4 | L | 600 | VR (2) | |
| <i>Ptilerodius pileatus</i> Capped Heron | O | 1 | L | 400 | U | |
| <i>Ardea cocoi</i> Cocoi Heron | O | 1 | L | 350 | U | |
| <i>Ardea alba</i> Great Egret | O | 1 | L | 350 | F | |
| <i>Bulbucius ibis</i> Cattle Egret | O | 1 | L | 400 | U | |
| <i>Egretta thula</i> Snowy Egret | O | 1 | L | 350 | F | |
| <i>Egretta caerulea</i> Little Blue Heron | O | 2 | 400 | 400 | VR (1) | |
| <i>Butorides striatus</i> Striated Heron | O | 1 | 350 | 350 | VR (1) | |
| <i>Agamia agami</i> Chestnut-bellied Heron | O | 2 | 350 | 350 | VR (1) | |

Appendix 2: Continued.

| Species | Evidence | Sites | Min | Max | Abundance | Breeding Information |
|--|----------|---------|-----|-------|-----------|----------------------|
| <i>Mesembrinitis cayennensis</i> Green Ibis | O | 1 | L | 400 | R | |
| <i>Mycteria americana</i> Wood Stork | O | 1 | L | 350 | VR (1) | |
| <i>Jaribu mycteria</i> Jabiru | O | 1 | L | 350 | VR (1) | |
| <i>Coragyps atratus</i> Black Vulture | O | 1,3-5 | L | 500 | C | |
| <i>Cathartes aura</i> Turkey Vulture | O | 1,3,5,6 | L | 900 | C | |
| <i>Cathartes burroivianus</i> Lesser Yellow-headed Vulture | O | 1 | L | 350 | R | |
| <i>Cathartes melambrotus</i> Greater Yellow-headed Vulture | O | 1-3,6 | L | 350 | R | |
| <i>Sarcorampus papa</i> King Vulture | O | 1,4,5 | L | 600 | U | |
| <i>Chauna torquata</i> Southern Screamer | O | 1 | L | 350 | VR (2) | |
| <i>Neochen jubata</i> Orinoco Goose | O | 1 | 350 | 350 | VR (1) | |
| <i>Cairina moschata</i> Muscovy Duck | O | 1,6 | L | 600 | VR (3) | |
| <i>Merganetta armata</i> Torrent Duck | O | 6 | 700 | 700 | VR (1Mfm) | |
| <i>Pandion haliaetus</i> Osprey | O | 1 | L | 400 | VR (1Mfm) | |
| <i>Chondrohierax uncinatus</i> Hook-billed Kite | O | 1,3-6 | L | 900 | U | |
| <i>Elanoides forficatus</i> Swallow-tailed Kite | O | 1,3,5,6 | L | 1,400 | F (+Mfm) | |
| <i>Harpagus bidentatus</i> Double-toothed Kite | T | 1,3,5 | L | 800 | R | |
| <i>Ictinia plumbea</i> Plumbeous Kite | O | 4,6 | L | 1,400 | VR (2) | |
| <i>Accipiter superciliosus</i> Tiny Hawk | T | 1,3,5,6 | L | 1,250 | VR (4) | F 10-10-96 |
| <i>Accipiter bicolor</i> Bicolored Hawk | O | 1 | L | 350 | VR (1) | |
| <i>Leucopternis albicollis</i> White Hawk | O | 1,3-5 | L | 700 | U | |
| <i>Buteogallus urubitinga</i> Great Black-hawk | O | 1,6 | L | 600 | VR (3) | |
| <i>Harpohaliaetus solitarius</i> Solitary Eagle | O | 5 | L | 900 | VR (2) | |
| <i>Buteo brachyurus</i> Short-tailed Hawk | O | 1,3,5 | L | 800 | F | |
| <i>Buteo albicaudatus</i> White-tailed Hawk | O | 5 | L | 700 | R | |
| <i>Harpia harpyja</i> Harpy Eagle | L | 1,6 | L | 600 | VR (2) | |
| <i>Spizaetus tyrannus</i> Black Hawk Eagle | O | 5 | 800 | 800 | VR (?) | |
| <i>Spizaetus ornatus</i> Ornate Hawk Eagle | O | 1,5 | L | 1,000 | VR (1) | |
| <i>Daptrius ater</i> Black Caracara | O | 1-2 | L | 500 | VR (2) | |
| <i>Ibycter americanus</i> Red-throated Caracara | T | 1-4,6 | L | 700 | F | |
| <i>Herpetotheres cachimans</i> Laughing Falcon | A | 1,4 | L | 500 | U | |

Appendix 2: Continued.

| Species | Evidence | Sites | Min | Max | Abundance | Breeding Information |
|---|----------|---------|-------|-------|-----------|----------------------|
| <i>Micrastur ruficollis</i> Barred Forest Falcon | T | 1,4-6 | L | 1,350 | F | |
| <i>Micrastur gilvicolis</i> Lined Forest Falcon | T | 4-6 | 700 | 1,200 | U | |
| <i>Micrastur mirandollei</i> Slaty-backed Forest Falcon | T | 1,2,5 | L | 700 | U | |
| <i>Micrastur semitorquatus</i> Collared Forest Falcon | T | 1,3,5 | L | 900 | U | |
| <i>Falco rufigularis</i> Bat Falcon | O | 1,3,5 | L | 800 | U | |
| <i>Ortalis guttata</i> Speckled Chachalaca | O | 1,2,5 | L | 500 | C | |
| <i>Penelope jacquacu</i> Spix's Guan | T | 1-6 | L | 1,300 | F | |
| <i>Pipile pipile</i> Blue-throated Piping Guan | O | 1,2,4-6 | L | 800 | U | |
| <i>Mitu tuberosa</i> Razor-billed Curassow | T | 1-6 | L | 1,000 | F | |
| <i>Odontophorus gujanensis</i> Marbled Wood Quail | T | 5-6 | L | 600 | U | |
| <i>Odontophorus speciosus</i> Rufous-breasted Wood Quail | T | 6 | L | 1,000 | R | |
| <i>Odontophorus balliviani</i> Stripe-faced Wood Quail * ** | T | 6 | 1,300 | 1,400 | U | F 15-09-98 |
| <i>Odontophorus stellatus</i> Starred Wood Quail | T | 1 | L | ?L | VR (1) | |
| <i>Aramidides cajana</i> Grey-necked Wood Rail | O | 1,2,5,6 | L | 700 | F | |
| <i>Eurypyga helias</i> Sunbittern | O | 1,2,5 | L | 800 | R | |
| <i>Psophia leucoptera</i> Pale-winged Trumpeter * | T | 2,4-6 | 500 | 1,000 | R | |
| <i>Vanellus cayanus</i> Pied Lapwing | O | 1 | L | 500 | U | |
| <i>Charadrius collaris</i> Collared Plover | O | 1,2 | L | 500 | F | |
| <i>Actitis macularia</i> Spotted Sandpiper | O | 1,2,5 | L | 500 | F | |
| <i>Bartramia longicauda</i> Upland Sandpiper | O | 6 | 1,300 | 1,300 | VR (1Mn) | |
| <i>Phaetusa simplex</i> Large-billed Tern | O | 1 | L | 500 | U | |
| <i>Sterna superciliaris</i> Yellow-billed Tern | O | 1 | L | 500 | U | |
| <i>Rynchops niger</i> Black Skimmer | O | 1 | L | 500 | U | |
| <i>Columba speciosa</i> Scaled Pigeon | T | 1-6 | L | 1,400 | U | |
| <i>Columba plumbea</i> Plumbeous Pigeon | T | 1-6 | L | 1,400 | C | |
| <i>Columba subvinacea</i> Ruddy Pigeon | O | 1,2,5 | L | 800 | U | |
| <i>Columbina picui</i> Picui Ground Dove | O | 1,5 | L | 500 | C | |
| <i>Columbina talpacoti</i> Ruddy Ground Dove | O | 1,5 | L | 500 | C | |
| <i>Claravis pretiosa</i> Blue Ground Dove | T | 1,3,6 | L | 500 | C | |
| <i>Claravis mondetoura</i> Maroon-chested Ground Dove | O | 3 | 800 | 800 | VR (1) | |
| <i>Leptotila verreauxi</i> White-tipped Dove | T | 2,5 | L | 600 | U | |

Appendix 2: Continued.

| Species | Evidence | Sites | Min | Max | Abundance | Breeding Information |
|---|----------|---------|-------|-------|-----------|----------------------|
| <i>Leptotila rufaxilla</i> Grey fronted Dove | T | 1-6 | L | 1,000 | C | |
| <i>Geotrygon frenata</i> White-throated Quail Dove | T | 6 | 1,000 | 1,400 | F | |
| <i>Geotrygon montana</i> Ruddy Quail Dove | O | 1,2,6 | L | 700 | R | |
| <i>Ara aurarana</i> Blue-and-yellow Macaw | O | 1-4 | L | 600 | F | |
| <i>Ara militaris</i> Military Macaw | T | 5,6 | 700 | 1,400 | U-R | |
| <i>Ara chloroptera</i> Red-and-green Macaw | T | 1-6 | L | 900 | F | NA 08-10-96 |
| <i>Ara scvera</i> Chestnut-fronted Macaw | O | 1-3,5 | L | 600 | C | |
| <i>Ara manilata</i> Red-bellied Macaw | O | 1,2 | L | 900 | R | |
| <i>Aratinga leucophthalmus</i> White-eyed Parakeet | O | 1-6 | L | 700 | C | |
| <i>Aratinga mitrata</i> Mitred Parakeet | T | 4,5 | 900 | 1,400 | F | |
| <i>Aratinga weddellii</i> Dusky-headed Parakeet | O | 1-6 | L | 600 | C | |
| <i>Pyrrhura molinae</i> Green-cheeked Parakeet | T | 6 | 1,200 | 1,400 | F | |
| <i>Pyrrhura picta</i> Painted Parakeet | T | 1,4,5 | L | 900 | U | |
| <i>Brotogeris cyanoptera</i> Cobalt-winged Parakeet | O | 1-3 | L | 600 | F | |
| <i>Pionites leucogaster</i> White-bellied Parrot * | O | 1,2 | L | 400 | U | |
| <i>Pionopsitta barrabandi</i> Orange-cheeked Parrot | O | 1 | L | 500 | VR (1) | |
| <i>Pionus menstruus</i> Blue-headed Parrot | T | 1-6 | L | 1,000 | F | |
| <i>Pionus sordidus</i> Red-billed Parrot | O | 5 | L | 800 | U | |
| <i>Amazona farinosa</i> Mealy Parrot | T | 1-6 | L | 1,000 | C | |
| <i>Piaya cayana</i> Squirrel Cuckoo | T | 1-6 | L | 1,400 | F | |
| <i>Crotophaga ani</i> Smooth-billed Ani | O | 1,4,5 | L | 500 | C | |
| <i>Tapera naevia</i> Striped Cuckoo | O | 1 | 450 | 450 | VR (1) | |
| <i>Dromococcyx phasianellus</i> Pheasant Cuckoo | O | 1 | L | 450 | R | |
| <i>Dromococcyx prazoninus</i> Pavonine Cuckoo | O | 1,5 | L | 500 | R | |
| <i>Neomorphus geoffroyi</i> Rufous-vented Ground Cuckoo | O | 5 | 700 | 700 | VR (1) | |
| <i>Otus guatemalae</i> Vermiculated Screech Owl | T | 4-6 | 600 | 800 | F | |
| <i>Otus choliba</i> Tropical Screech Owl | O | 1,2 | L | 500 | F | |
| <i>Otus ingens</i> Rufescent Screech Owl | T | 1,4-6 | 900 | 1,300 | F | |
| <i>Otus watsonii</i> Tawny-bellied Screech Owl | T | 1,2,5,6 | L | 900 | F | |
| <i>Lophostrix cristata</i> Crested Owl | O | 1-6 | L | 600 | F | |
| <i>Pulsatrix perspicillata</i> Spectacled Owl | O | 1,3,5 | L | 600 | F | |

Appendix 2: Continued.

| Species | Evidence | Sites | Min | Max | Abundance | Breeding Information |
|--|----------|---------|-----|-------|------------|----------------------|
| <i>Pulsatrix melanota</i> Band-bellied Owl | T | 4-6 | 700 | 1,200 | F | |
| <i>Glaucidium hardyi</i> Amazonian Pygmy Owl | T | 1,2,5 | L | 700 | U | |
| <i>Glaucidium parkeri</i> Subtropical Pygmy Owl | T | 5,6 | 700 | 1,200 | U | |
| <i>Glaucidium brasilianum</i> Ferruginous Pygmy Owl | T | 1 | L | 600 | F | |
| <i>Ciccaba virgata</i> Mottled Owl | T | 1,5 | L | 700 | U | |
| <i>Ciccaba huluha</i> Black-banded Owl | T | 1,2,4-6 | L | 1,000 | F | |
| <i>Steatornis carpinensis</i> Oilbird | T | 6 | 400 | 400 | C | |
| <i>Nyctibius grandis</i> Great Potoo | T | 1,2,6 | L | 800 | R | |
| <i>Nyctibius aethereus</i> Long-tailed Potoo | T | 1,5 | L | 900 | R | |
| <i>Nyctibius griseus</i> Common Potoo | O | 1 | L | 500 | R | |
| <i>Chordeiles rupestris</i> Sand-colored Nighthawk | O | 1 | L | 500 | R | |
| <i>Nyctidromus albigollis</i> Parouque | O | 1,4 | L | 500 | R | |
| <i>Nyctiphrynus ocellatus</i> Ocellated Poorwill | T | 5 | L | 900 | R | |
| <i>Caprimulgus rufus</i> Rufous Nighthawk | O | 5 | L | 900 | VR (2)M(s) | |
| <i>Caprimulgus serripinnatus</i> Silky-tailed Nighthawk | O | 1 | L | 500 | VR (1) | |
| <i>Hydrophalis climacocera</i> Ladder-tailed Nighthawk | SP | 1 | L | 500 | VR (2) | |
| <i>Streptoprocne rutila</i> Chestnut-collared Swift | O | 1 | L | 500 | VR (1) | |
| <i>Chaetura cinereiventris</i> Grey-rumped Swift | O | 1-6 | L | 1,400 | C | |
| <i>Chaetura brachyura</i> Short-tailed Swift | O | 1,5 | L | 900 | U | |
| <i>Threnetes leucurus</i> Pale-tailed Barbthroat | O | 1,5 | L | 1,400 | F-U | |
| <i>Phaethornis malaris</i> Long-tailed Hermit | T | 1,4,5 | L | 900 | R | |
| <i>Phaethornis hispidus</i> White-bearded Hermit | O | 1,4-6 | L | 1,300 | F | |
| <i>Phaethornis stuarti</i> White-browed Hermit * | O | 1,3-6 | L | 800 | U | |
| <i>Phaethornis ruber</i> Reddish Hermit | T | 5 | 700 | 700 | VR (1) | |
| <i>Campylopterus largipennis</i> Grey-breasted Sabrewing | O | 1,3-6 | L | 1,400 | C | |
| <i>Florisuga mellivora</i> White-necked Jacobin | O | 1,3-6 | L | 1,300 | U | |
| <i>Anthracoceros nigricollis</i> Black-throated Mango | O | 1,5-6 | L | 850 | U | |
| <i>Klais guimeti</i> Violet-headed Hummingbird | O | 1,6 | L | 900 | R | |
| <i>Lophornis delattrei</i> Rufous-crested Coquette | T | 3-5 | 700 | 1,000 | U | |
| <i>Lophornis chalybeus</i> Festive Coquette | O | 5 | 900 | 900 | VR (1) | |
| | O | 1,5 | L | 600 | F | |

Appendix 2: Continued.

| Species | Evidence | Sites | Min | Max | Abundance | Breeding Information |
|--|----------|---------|-------|-------|-----------|----------------------|
| <i>Chlorostilbon mellisugus</i> Blue-tailed Emerald | O | 1,6 | L | 1,200 | R | |
| <i>Thalassia furcata</i> Fork-tailed Woodnymph | O | 1,5-6 | L | 1,400 | U | |
| <i>Hylocharis cyanus</i> White-chinned Sapphire | O | 1,5 | L | 700 | R | |
| <i>Chrysoronia oenone</i> Golden-tailed Sapphire | T | 1,4-5 | L | 900 | F | |
| <i>Tapirospilus hypostictus</i> Many-spotted Hummingbird | O | 5 | 875 | 875 | VR (1) | |
| <i>Amazilia versicolor</i> Versicolored Emerald | O | 1,5 | L | 900 | U | |
| <i>Amazilia lactea</i> Sapphire-spangled Emerald | O | 5 | 875 | 875 | VR (1) | |
| <i>Adelomyia melanogenys</i> Speckled Hummingbird | O | 6 | 1,200 | 1,450 | F | |
| <i>Polyplancta aurescens</i> Gould's Jewelfront | T | 1,4-5 | L | 900 | U | |
| <i>Heliothryx aurita</i> Black-eared Fairy | O | 5,6 | 875 | 900 | VR (2) | |
| <i>Calliphlox amethystina</i> Amethyst Woodstar | O | 1,5 | L | 800 | U | |
| <i>Pharomacrus antisianus</i> Crested Quetzal | T | 4,6 | 900 | 1,400 | F | |
| <i>Trogon melanurus</i> Black-tailed Trogon | T | 1-6 | L | 1,000 | F | |
| <i>Trogon viridis</i> White-tailed Trogon | T | 2,5,6 | L | 700 | U | |
| <i>Trogon collaris</i> Collared Trogon | T | 1,2,4,5 | L | 1,200 | U | |
| <i>Trogon curucui</i> Blue-crowned Trogon | T | 1-6 | L | 1,400 | F | |
| <i>Trogon violaceus</i> Violaceous Trogon | O | 1 | L | 450 | U | |
| <i>Electron platyrhynchum</i> Broad-billed Motmot | O | 1-6 | L | 1,000 | U | |
| <i>Baryphthengus martii</i> Rufous Motmot | T | 1-3,5,6 | 650 | 1,000 | F-U | |
| <i>Momotus momota</i> Blue-crowned Motmot | T | 1-3,5 | L | 500 | F | |
| <i>Ceryle torquata</i> Ringed Kingfisher | O | 1,2,4,5 | L | 600 | F | |
| <i>Chloroceryle amazona</i> Amazon Kingfisher | T | 1,2,5 | L | 500 | U | |
| <i>Chloroceryle americana</i> Green Kingfisher | O | 1,5 | L | 500 | U | |
| <i>Chloroceryle inda</i> Green-and-rufous Kingfisher | O | 2 | L | 500 | R | |
| <i>Chloroceryle aenea</i> Pygmy Kingfisher | O | 1,6 | L | 500 | R | |
| <i>Notiarchus macrotympanus</i> White-necked Puffbird | T | 4-6 | 600 | 900 | U | |
| <i>Bucco macrodactylus</i> Chestnut-capped Puffbird | O | 1 | L | 450 | VR (1) | |
| <i>Nyctialus striolatus</i> Striolated Puffbird | T | 1,3-6 | L | 800 | U | |
| <i>Micromonacha lanceolata</i> Lanceolated Monklet | O | 6 | 1,150 | 1,200 | VR (2) | |
| <i>Monasa nigrifrons</i> Black-fronted Nunbird | O | 1,2 | L | 500 | F | |
| <i>Monasa morphoeus</i> White-fronted Nunbird | O | 1,2,5,6 | L | 500 | F | |

Appendix 2: Continued.

| Species | Evidence | Sites | Min | Max | Abundance | Breeding Information |
|--|----------|---------|-------|-------|-----------|----------------------|
| <i>Chelidoptera tenebrosa</i> Swallow-wing | O | 1,4,6 | L | 500 | U | |
| <i>Galbula ruficauda</i> Rufous-tailed Jacamar | O | 1,2,5,6 | L | 500 | F | |
| <i>Capito niger</i> Black-spotted Barbet | T | 1-6 | L | 950 | C | |
| <i>Eubucco richardsoni</i> Lemon-throated Barbet | O | 1,4-6 | L | 850 | R | |
| <i>Eubucco versicolor</i> Versicolored Barbet * | T | 6 | 1,200 | 1,400 | F | |
| <i>Aulacorhynchus prasinus</i> Emerald Toucanet | T | 4-6 | L | 1,000 | F | |
| <i>Aulacorhynchus derbianus</i> Chestnut-tipped Toucanet | T | 5,6 | 800 | 1,400 | F | |
| <i>Pteroglossus inscriptus</i> Lettered Aracari | O | 6 | 800 | 800 | VR (1) | |
| <i>Pteroglossus flavirostris</i> Ivory-billed Aracari | O | 1,5,6 | L | 1,000 | U | |
| <i>Pteroglossus castanotis</i> Chestnut-eared Aracari | T | 1,2,5,6 | L | 1,000 | C | |
| <i>Pteroglossus beauharnaisii</i> Curl-crested Aracari * | T | 1,5 | L | 800 | R | |
| <i>Selenidera reinwardtii</i> Golden-collared Toucanet | T | 1,2,4-6 | L | 800 | R | |
| <i>Ramphastos vitellinus</i> Channel-billed Toucan | T | 1-6 | L | 1,400 | C | |
| <i>Ramphastos tucanus</i> Red-billed Toucan | T | 1-6 | L | 1,000 | C | |
| <i>Picumnus minutissimus</i> Arrowhead Piculet | O | 4,5 | L | 800 | R | |
| <i>Picumnus aurifrons</i> Bar-breasted Piculet * | T | 4,5 | L | 900 | R | |
| <i>Melanerpes cruentatus</i> Yellow-tufted Woodpecker | T | 1-6 | L | 700 | C | |
| <i>Ventilornis fumigatus</i> Smoky-brown Woodpecker | T | 6 | 1,300 | 1,400 | U | |
| <i>Ventilornis passerinus</i> Little Woodpecker | O | 1,3,4 | L | 800 | U | |
| <i>Ventilornis affinis</i> Red-stained Woodpecker | O | 1,5 | 400 | 1,000 | F | |
| <i>Piculus leucolaemus</i> White-throated Woodpecker | T | 1,3-6 | L | 1,000 | F | |
| <i>Piculus chrysocoloros</i> Golden-green Woodpecker | T | 5,6 | 500 | 700 | VR (3) | |
| <i>Piculus rubiginosus</i> Golden-olive Woodpecker | T | 5,6 | 500 | 1,400 | U | |
| <i>Ceuleus gramineus</i> Scale-breasted Woodpecker | T | 4 | 900 | 900 | VR (1) | |
| <i>Ceuleus flavus</i> Cream-coloured Woodpecker | O | 1 | L | 500 | R | |
| <i>Dryocopus lineatus</i> Lineated Woodpecker | O | 1,5 | 500 | 1,000 | R | |
| <i>Campophilus melanoleucos</i> Crimson-crested Woodpecker | O | 5 | 400 | 400 | VR (1) | |
| <i>Campophilus rubricollis</i> Red-necked Woodpecker | T | 1-6 | L | 1,300 | F | |
| <i>Dendrocincla fuliginosa</i> Plain-brown Woodcreeper | T | 1,2,4-6 | L | 1,000 | F | F 08-10-96 |
| <i>Deconychura longicauda</i> Long-tailed Woodcreeper | T | 1,5,6 | L | 900 | U | |
| <i>Sittasomus griseicapillus</i> Olivaceous Woodcreeper | T | 1,3-6 | L | 1,300 | F | |

Appendix 2: Continued.

| Species | Evidence | Sites | Min | Max | Abundance | Breeding Information |
|--|----------|---------|-------|-------|-----------|----------------------|
| <i>Dendrexetastes rufigula</i> Cinnamon-throated Woodcreeper | T | 4,5 | L | 800 | R | |
| <i>Xiphocolaptes promeropi</i> Xiphichus Strong-billed Woodcr. | T | 1,4-6 | L | 900 | F | |
| <i>Dendrocolaptes certhia</i> Banded Woodcreeper | T | 1,4 | L | 700 | F | |
| <i>Dendrocolaptes picumnus</i> Black-banded Woodcreeper | T | 4-6 | L | 900 | U | |
| <i>Xiphorhynchus picus</i> Straight-billed Woodcreeper | T | 1,6 | L | 800 | R | |
| <i>Xiphorhynchus ocellatus</i> Ocellated Woodcreeper | T | 1,3,5 | L | 900 | U | |
| <i>Xiphorhynchus spixii</i> Spix's Woodcreeper | T | 1-6 | L | 900 | F | |
| <i>Xiphorhynchus guttatus</i> Buff-throated Woodcreeper | T | 1-6 | L | 1,000 | F | |
| <i>Lepidocolaptes albolineatus</i> Lineated Woodcreeper | T | 1,5,6 | L | 900 | U | |
| <i>Campylorhamphus trochilirostris</i> Red-billed Scythebill | SP | 1,5,6 | L | 1,300 | U | |
| <i>Furnarius leucopus</i> Pale-legged Hornero | O | 1 | L | 350 | U | |
| <i>Synallaxis azarae</i> Azara's Spinetail * | T | 6 | 1,300 | 1,400 | F | |
| <i>Synallaxis cabanisi</i> Cabanis' Spinetail * | T | 5,6 | 800 | 1,300 | F | |
| <i>Synallaxis gujanensis</i> Plain-crowned Spinetail | T | 1,2 | L | 500 | F | |
| <i>Cranioleuca curtata</i> Ash-browed Spinetail | T | 5,6 | 700 | 900 | R | |
| <i>Cranioleuca gutturata</i> Speckled Spinetail | O | 1 | L | 500 | VR (1) | |
| <i>Prennoplex brunescens</i> Spotted Barbtail | T | 6 | 1,300 | 1,300 | VR (1) | |
| <i>Hylocistis subulatus</i> Striped Woodhaunter | O | 5,6 | 700 | 1,300 | VR (3) | |
| <i>Ancistrops strigilatus</i> Chestnut-winged Hookbill | O | 5 | 800 | 800 | VR (1) | |
| <i>Simoxenops striatus</i> Bolivian Recurvebill *, **, *** | T | 4,6 | 700 | 1,300 | U-R | |
| <i>Philydor erythrocerus</i> Rufous-rumped Foliage-gleaner | O | 3,4 | 600 | 900 | VR (3) | |
| <i>Philydor erythropterus</i> Chestnut-winged Foliage-gleaner | O | 5 | 700 | 850 | VR (2) | |
| <i>Automolus ochrolaemus</i> Buff-throated Foliage-gleaner | T | 1,2,4-6 | L | 1,400 | F | |
| <i>Xenops rutilans</i> Streaked Xenops | O | 1,5,6 | 500 | 700 | U | |
| <i>Xenops minutus</i> Plain Xenops | SP | 1,5,6 | 600 | 900 | U | |
| <i>Sclerurus albigularis</i> Grey-throated Leaf-tosser | T | 3,5,6 | 700 | 800 | R | |
| <i>Sclerurus mexicanus</i> Tawny-throated Leaf-tosser | T | 4,5 | 700 | 800 | U | |
| <i>Cymbilaimus lineatus</i> Fasciated Antshrike | O | 1,4,5 | L | 1,000 | F | |
| <i>Taraba major</i> Great Antshrike | O | 1,2,5,6 | L | 1,000 | C | |
| <i>Thamnophilus palliatus</i> Chestnut-backed Antshrike | T | 6 | 1,200 | 1,400 | R | |
| <i>Thamnophilus nethiops</i> White-shouldered Antshrike | T | 1,4-6 | L | 1,000 | F | |

Appendix 2: Continued.

| Species | Evidence | Sites | Min | Max | Abundance | Breeding Information |
|---|----------|---------|-------|-------|-----------|----------------------|
| <i>Thamnoptilus schistaceus</i> Plain-winged Antshrike | T | 1-6 | L | 1,000 | C | |
| <i>Thamnoptilus aroyae</i> Upland Antshrike*, ** | T | 5,6 | 1,200 | 1,400 | F | |
| <i>Thamnoptilus caeruleus</i> Variable Antshrike | T | 6 | 1,300 | 1,300 | VR (1) | |
| <i>Dysithamnus mentalis</i> Plain Antvireo | T | 1,3,5,6 | 450 | 1,000 | F | |
| <i>Thamnomantes schistogynus</i> Bluish-slate Antshrike * | T | 2,4,5 | 600 | 1,000 | U | |
| <i>Myrmotherula brachyura</i> Pygmy Antwren | T | 1,4,5 | L | 1,000 | F | |
| <i>Myrmotherula longicauda</i> Stripe-chested Antwren | O | 1,5 | 600 | 900 | R | |
| <i>Myrmotherula hauxwelli</i> Plain-throated Antwren | O | 1 | L | L | VR (1) | |
| <i>Myrmotherula leucophthalma</i> White-eyed Antwren * | O | 5,6 | L | 900 | VR (2) | |
| <i>Myrmotherula axillaris</i> White-flanked Antwren | O | 1,5,6 | L | 1,000 | U | |
| <i>Myrmotherula grisea</i> Yungas Antwren *, **, *** | O | 5 | 850 | 850 | VR (1) | |
| <i>Myrmotherula menetriesii</i> Grey Antwren | T | 4,5 | L | 1,000 | F | |
| <i>Dichrozona cincta</i> Banded Antbird | T | 4,5 | 500 | 800 | U | |
| <i>Herpilochmus rufimarginatus</i> Rufous-winged Antwren | T | 2-6 | L | 1,000 | C-F | |
| <i>Drymophila devillei</i> Striated Antbird | T | 2,5,6 | 700 | 1,350 | F | |
| <i>Cercomacra cinerascens</i> Grey Antbird | T | 1,2,4-6 | L | 1,000 | C | |
| <i>Cercomacra nigrescens</i> Blackish Antbird | O | 1 | L | L | R | NM 07-10-98 |
| <i>Cercomacra serna</i> Black Antbird | T | 1,4-6 | L | 1,000 | U | |
| <i>Pyriglena leuconota</i> White-backed Fire-eye | T | 5,6 | 700 | 1,400 | F | |
| <i>Myrmoborus leucophrys</i> White-browed Antbird | O | 2 | L | L | R | |
| <i>Myrmoborus myotherinus</i> Black-faced Antbird | T | 1-6 | L | 1,400 | C | |
| <i>Hypocnemis cantator</i> Warbling Antbird | T | 2,5,6 | L | 800 | C | |
| <i>Percnostola leucostigma</i> Spot-winged Antbird | O | 1 | L | L | VR (2) | |
| <i>Myrmeciza hemimelaena</i> Chestnut-tailed Antbird | T | 1,2,4-6 | L | 1,200 | C | NE 21-09-97 |
| <i>Myrmeciza atrothorax</i> Black-throated Antbird | T | 1,2,4-6 | L | 1,000 | C | |
| <i>Rhegnatorhina melanosticta</i> Hairy-crested Antbird | T | 4-6 | 500 | 900 | R | |
| <i>Hylophylax naevia</i> Spot-backed Antbird | SP | 1,2,4-6 | L | 1,000 | U | |
| <i>Hylophylax poecilnota</i> Scale-backed Antbird | O | 5 | 600 | 600 | VR (1) | |
| <i>Formicarius colma</i> Rufous-capped Antthrush | T | 1 | L | 500 | R | |
| <i>Formicarius analis</i> Black-faced Antthrush | T | 1-6 | L | 1,000 | C | |
| <i>Chamaeza campanisona</i> Short-tailed Antthrush | T | 1,5,6 | 1,200 | 1,400 | C | |

Appendix 2: Continued.

| Species | Evidence | Sites | Min | Max | Abundance | Breeding Information |
|---|----------|---------|-------|-------|-----------|----------------------|
| <i>Gnallaria guatemalensis</i> Scaled Antpitta | T | 5,6 | 700 | 1,000 | R | |
| <i>Hylopezus berlepschi</i> Amazonian Antpitta * | O | 1 | 350 | 350 | VR (1) | |
| <i>Conopophaga ardesiaca</i> Slaty Gnateater * | T | 6 | 1,200 | 1,400 | U | |
| <i>Scytalopus bolivianus</i> White-crowned Tapaculo | T | 6 | 1,200 | 1,400 | C | |
| <i>Phyllomyias burmeisteri</i> Rough-legged Tyrannulet | O | 5 | 800 | 1,000 | U? | |
| <i>Phyllomyias sclateri</i> Sclater's Tyrannulet * | O | 5 | 480 | 1,200 | F | |
| <i>Phyllomyias</i> , undescribed species * **, *** | T | 1,5 | 875 | 1,000 | U? | |
| <i>Zimmerius gracilipes</i> Slender-footed Tyrannulet | T | 1,5,6 | L | 1,100 | C | |
| <i>Ornithion inerne</i> White-lored Tyrannulet | O | 1,3-5 | L | 900 | U | |
| <i>Camplostoma obsoletum</i> S. Beardless Tyrannulet | O | 5 | 800 | 800 | VR (1Ms) | |
| <i>Tyrannulus elatus</i> Yellow-crowned Tyrannulet | O | 1 | L | 500 | VR (1) | |
| <i>Myiopagis gainardii</i> Forest Elaenia | T | 1,5 | L | 950 | F | |
| <i>Myiopagis caniceps</i> Grey Elaenia | O | 4,5 | L | 900 | R | |
| <i>Myiopagis viridicata</i> Greenish Elaenia | O | 1 | L | 500 | ? | |
| <i>Elaenia albiceps</i> White-crested Elaenia | T | 5 | L | 1,000 | R | |
| <i>Elaenia obscura</i> Highland Elaenia | O | 4,5 | 800 | 950 | R | |
| <i>Serpophaga cinerea</i> Torrent Tyrannulet | O | 6 | 1,100 | 1,100 | VR (1Mm) | |
| <i>Mionectes striatocollis</i> Streak-necked Flycatcher | T | 6 | 1,350 | 1,350 | VR (1Mm) | |
| <i>Mionectes oleagineus</i> Ochre-bellied Flycatcher | T | 1,2,5,6 | L | 1,100 | F | F 15-11-99 |
| <i>Mionectes macconnelli</i> McConnell's Flycatcher | T | 1,4-6 | L | 1,000 | U | |
| <i>Leptopogon amaurocephalus</i> Sepia-capped Flycatcher | T | 1,5,6 | L | 1,000 | F | |
| <i>Leptopogon superciliosus</i> Slaty-capped Flycatcher | T | 5,6 | 850 | 1,300 | U | |
| <i>Phylloscartes ophthalmicus</i> Marble-faced Bristle Tyrant | T | 5 | 800 | 1,000 | R | |
| <i>Phylloscartes orbitalis</i> Spectacled Bristle Tyrant | O | 5,6 | 700 | 1,000 | U | |
| <i>Phylloscartes flaviventris</i> Rufous-lored Tyrannulet | O | 5 | 900 | 900 | VR (1) | |
| <i>Corythopis torquata</i> Ringed Antpitt | T | 2,5,6 | L | 950 | U | |
| <i>Myiornis albiventris</i> White-bellied Pygmy Tyrant * | T | 2,4-6 | 600 | 1,100 | F | |
| <i>Myiornis ecaudatus</i> Short-tailed Pygmy Tyrant | T | 1,3-6 | L | 950 | F | |
| <i>Hemitriccus flammulatus</i> Flammulated Tody Tyrant * | O | 5 | L | 700 | R | |
| <i>Hemitriccus zosterops</i> White-eyed Tody Tyrant | T | 1,2,5,6 | L | 900 | F | |
| <i>Hemitriccus spodiops</i> Yungas Tody Tyrant * **, *** | T | 5,6 | 1,200 | 1,400 | C | NM 27-07-97 |

Appendix 2: Continued.

| Species | Evidence | Sites | Min | Max | Abundance | Breeding Information |
|------------------------------------|----------|-------|-------|-------|-----------|----------------------|
| <i>Hemitriccus rufigularis</i> | T | 5,6 | 800 | 1,350 | R | |
| <i>Todirostrum plumbeiceps</i> | O | 5 | L | 500 | U | |
| <i>Todirostrum chrysocrotaphum</i> | O | 1,5,6 | 0 | 1,000 | R | |
| <i>Ramphotrigon ruficauda</i> | O | 1 | 0 | 600 | VR (1) | |
| <i>Rhynchocyclus olivaceus</i> | T | 1,5 | 0 | 950 | R | |
| <i>Tolmomyias sulphureus</i> | T | 1,5,6 | L | 900 | R | |
| <i>Tolmomyias poliocephalus</i> | T | 1,5 | L | ? | ? | NM 02-08-97 |
| <i>Tolmomyias flaviventris</i> | T | 4 | 800 | 800 | VR (2) | |
| <i>Platyrinchus coronatus</i> | O | 1,2 | L | 600 | R | |
| <i>Terenotriccus erythurus</i> | O | 5 | 800 | 1,100 | R | |
| <i>Myiobius villosus</i> | T | 6 | 1,200 | 1,350 | R | |
| <i>Myiophobus fasciatus</i> | T | 1,5 | L | 1,000 | F | |
| <i>Pyrrhomyias cinnamomea</i> | O | 6 | 900 | 900 | VR (1Mm) | |
| <i>Contopus fumigatus</i> | O | 5 | 500 | 1,000 | R | |
| <i>Contopus virens</i> | T | 2-5 | L | 1,000 | C-F (Mm) | |
| <i>Contopus cinereus</i> | O | 5 | L | 1,000 | R | |
| <i>Empidonax euleri</i> | T | 1,4-6 | L | 1,000 | F | |
| <i>Cnemotriccus fuscatus</i> | T | 5 | 600 | 600 | VR (1) | |
| <i>Sayornis nigricans</i> | O | 5,6 | L | 900 | U | |
| <i>Pyrocephalus rubinus</i> | O | 1 | L | 500 | F (Ms) | |
| <i>Ochthoeca littoralis</i> | O | 1,2,5 | L | 500 | F | |
| <i>Muscisaxicola fluvialis</i> | O | 1 | L | 500 | F | |
| <i>Knipolegus signatus</i> | O | 5 | 1,200 | 1,200 | VR (1Mm) | |
| <i>Colonia colonus</i> | O | 6 | L | 500 | R | |
| <i>Satrapa icterophrys</i> | O | 5 | 440 | 440 | VR (1) | |
| <i>Attila bolivianus</i> | T | 3, 4 | 450 | 700 | U | |
| <i>Attila spadiceus</i> | T | 1,3-6 | L | 1,300 | F | F 21-11-99 |
| <i>Rhytipterna simplex</i> | T | 1-6 | L | 1,000 | F-U | |
| <i>Sirystes sibilator</i> | O | 5 | L | 700 | R | |
| <i>Myiarchus tuberculifer</i> | T | 1 | L | 500 | R | |
| <i>Myiarchus ferox</i> | T | 1,5 | L | 700 | F | |

Appendix 2: Continued.

| Species | Evidence | Sites | Min | Max | Abundance | Breeding Information |
|--|----------|---------|-----|-------|-----------|----------------------|
| <i>Mniotilta tyrannuloides</i> Brown-crested Flycatcher | T | 1,5 | L | 900 | U | |
| <i>Pitangus sulphuratus</i> Great Kiskadee | O | 1 | L | 500 | F | |
| <i>Megarynchus pitangua</i> Boat-billed Flycatcher | O | 1,5 | L | 600 | F | |
| <i>Myiozetetes cayanensis</i> Rusty-margined Flycatcher | T | 1,5 | 450 | 800 | R | |
| <i>Myiozetetes similis</i> Social Flycatcher | O | 1,5 | L | 700 | R | |
| <i>Myiozetetes luteiventris</i> Dusky-chested Flycatcher | O | 1 | L | 500 | VR (1) | |
| <i>Conopias trivirgata</i> Three-striped Flycatcher | O | 6 | L | 900 | R | |
| <i>Myiodinastes chryscephalus</i> Golden-crowned Flycatcher | T | 1,5 | L | 1,400 | C-F | |
| <i>Myiodinastes maculatus</i> Streaked Flycatcher | T | 1-3,5,6 | L | 1,000 | F | |
| <i>Myiodinastes luteiventris</i> Sulphur-bellied Flycatcher | T | 4-5 | 500 | 900 | F (Mn) | |
| <i>Legatus leucophaeus</i> Piratic Flycatcher | T | 1-6 | L | 1,000 | F | |
| <i>Empidonax varius</i> Variegated Flycatcher | O | 3 | 600 | 600 | VR (1) | NA 12-10-96 |
| <i>Empidonax aurantiotrochistatus</i> Crowned Slaty Flycatcher | O | 5 | 700 | 700 | VR (1) | |
| <i>Tyrannus albogularis</i> White-throated Kingbird | O | 5,6 | 700 | 950 | R | |
| <i>Tyrannus melancholicus</i> Tropical Kingbird | T | 1,3-6 | L | 1,200 | C | |
| <i>Tyrannus tyrannus</i> Eastern Kingbird | O | 1,4 | L | 500 | R (Mn) | |
| <i>Pachyrhamphus castaneus</i> Chestnut-crowned Becard | O | 1 | L | L | VR (1) | |
| <i>Pachyrhamphus polychopterus</i> White-winged Becard | O | 1,4,5 | L | 800 | R | |
| <i>Pachyrhamphus marginatus</i> Black-capped Becard | O | 1,4-6 | L | 1,000 | R | |
| <i>Pachyrhamphus minor</i> Pink-throated Becard | T | 1,3-5 | L | 700 | U | |
| <i>Pachyrhamphus validus</i> Crested Becard | O | 1,4 | L | 1,000 | VR (2) | NB 08-10-96 |
| <i>Tityra cayana</i> Black-tailed Tityra | O | 1 | L | L | VR (1) | |
| <i>Tityra semifasciata</i> Masked Tityra | T | 1-6 | L | 1,300 | F | NM 27-07-97 |
| <i>Laniusoma elegans</i> Shrike-like Cotinga | O | 6 | 850 | 850 | VR (1) | |
| <i>Amphispiza bilineata</i> Scaled Fruiteater | T | 4 | 850 | 900 | R | |
| <i>Laniocera hypopyrra</i> Cinereous Mourner | O | 1,4 | L | 500 | R | |
| <i>Lipaugus vociferans</i> Screaming Piha | T | 1-6 | L | 900 | C | |
| <i>Cotinga cayana</i> Spangled Cotinga | O | 1,6 | L | 800 | VR (2) | |
| <i>Gymnoderus foetidus</i> Bare-necked Fruitcrow | O | 1,6 | L | 600 | VR (2) | |
| <i>Querula purpurata</i> Purple-throated Fruitcrow | T | 1,3-6 | L | 900 | F | |
| <i>Cephalopterus ornatus</i> Amazonian Umbrellabird | O | 6 | 900 | 900 | VR (1) | |

Appendix 2: Continued.

| Species | Evidence | Sites | Min | Max | Abundance | Breeding Information |
|------------------------------------|----------|---------|-------|-------|-----------|----------------------|
| <i>Rupicola peruviana</i> | T | 6 | 700 | 1,200 | U | |
| <i>Schiffornis turdinus</i> | T | 1,2,4-6 | 800 | 1,400 | F | F 06-10-98 |
| <i>Piprites chloris</i> | T | 1-6 | 600 | 1,000 | F | |
| <i>Tyranneteutes stolzmanni</i> | O | 1,2,6 | L | 800 | R | |
| <i>Machaeropterus pyrocephalus</i> | SP | 1,5 | L | 400 | VR (2) | |
| <i>Chiroxipha boliviana</i> | T | 5,6 | 1,000 | 1,400 | C | |
| <i>Pipra coronata</i> | O | 2,6 | 600 | 1,200 | R | |
| <i>Pipra fasciicauda</i> | SP | 1,2 | L | 500 | R | |
| <i>Pipra chloromeros</i> | T | 1-6 | L | 800 | C | |
| <i>Tachycineta albiventer</i> | O | 1 | L | 500 | C | |
| <i>Progne chalybea</i> | O | 1 | 400 | 400 | VR (1) | |
| <i>Notiochelidon cyanoleuca</i> | T | 3,5 | 700 | 1,400 | F | |
| <i>Aticora fasciata</i> | O | 1,2,4 | L | 500 | C | |
| <i>Stelgidopteryx ruficollis</i> | O | 1-5 | L | 900 | C | |
| <i>Campylorhynchus turdinus</i> | T | 1,3-5 | L | 500 | F | |
| <i>Thryothorus genibarbis</i> | T | 1-6 | L | 800 | C | |
| <i>Thryothorus guarayanus</i> | T | 1 | 500 | 500 | VR (1) | |
| <i>Troglodytes aedon</i> | T | 1,3,5 | L | 1,300 | F | |
| <i>Henicorhina leucophrys</i> | T | 6 | 1,200 | 1,400 | C | |
| <i>Microcerculus marginatus</i> | T | 1-6 | L | 900 | C | |
| <i>Cyphorhinus arada</i> | T | 1 | L | 400 | U | |
| <i>Donacobius atricapillus</i> | O | 1,5 | L | 500 | F | |
| <i>Cinclus leucocephalus</i> | O | 6 | 700 | 700 | VR (1Mm) | |
| <i>Ramphocaeus melanurus</i> | T | 1-5 | L | 900 | F | |
| <i>Catharus ustulatus</i> | T | 4-6 | 700 | 1,000 | F (Mn) | |
| <i>Platycichla leucops</i> | T | 6 | 1,400 | 1,400 | VR (1) | |
| <i>Turdus nigricaps</i> | O | 5 | L | 700 | VR (2) | |
| <i>Turdus amaurochalinus</i> | T | 1,3-5 | L | 700 | F | |
| <i>Turdus ignobilis</i> | O | 1,3-5 | L | 1,000 | F | |
| <i>Turdus lawrencii</i> | O | 1,2 | L | 500 | U | |
| <i>Turdus hauxwelli</i> | O | 1 | L | L | R | |

Appendix 2: Continued.

| Species | Evidence | Sites | Min | Max | Abundance | Breeding Information |
|--|----------|---------|-------|-------|-----------|----------------------|
| <i>Turdus albicollis</i> White-necked Thrush | O | 5 | L | 900 | F | |
| <i>Cyclarhis guianensis</i> Rufous-browed Peppershrike | T | 1,3 | L | 500 | U | |
| <i>Vireolanus leucotis</i> Slaty-capped Shrike-Vireo | T | 1,3-6 | L | 900 | U | |
| <i>Vireo olivaceus</i> Red-eyed Vireo | T | 1-6 | L | 1,000 | F | |
| <i>Vireo flavoviridis</i> Yellow-green Vireo | O | 5 | 700 | 900 | R (Mn) | |
| <i>Vireo leucophrys</i> Brown-capped Vireo | O | 5 | 600 | 1,000 | R | |
| <i>Hylophilus thoracicus</i> Lemon-chested Greenlet | O | 5,6 | 600 | 1,000 | R | |
| <i>Hylophilus hypoxanthus</i> Dusky-capped Greenlet | T | 1-6 | L | 900 | C | |
| <i>Hylophilus ochraceiceps</i> Tawny-crowned Greenlet | T | 1-6 | L | 900 | C | |
| <i>Ammodramus aurifrons</i> Yellow-browed Sparrow | O | 1,2,4-6 | L | 600 | F | |
| <i>Cyanocorax cyanomelas</i> Purplish Jay | O | 1,2,5 | L | 700 | C | |
| <i>Cyanocorax yncas</i> Green Jay | T | 5,6 | 1,200 | 1,400 | U | |
| <i>Volatinia jacarina</i> Blue-black Grassquit | O | 1 | L | L | VR (1Ms) | |
| <i>Sporophila collaris</i> Rusty-collared Seedeater | O | 5 | L | L | VR (1Ms) | |
| <i>Sporophila lineola</i> Lined Seedeater * | O | 1 | L | L | VR (1Ms) | |
| <i>Sporophila caerulescens</i> Double-collared Seedeater | O | 1,2,3 | L | L | R (Ms) | |
| <i>Sporophila castaneiventris</i> Chestnut-bellied Seedeater | O | 1 | L | L | VR (2Ms) | |
| <i>Oryzoborus angolensis</i> Lesser Seedfinch | O | 5 | L | 700 | R | |
| <i>Arremon taciurnus</i> Pectoral Sparrow | T | 1-6 | L | 1,000 | C | |
| <i>Paroaria gularis</i> Red-capped Cardinal | O | 1 | 500 | 500 | VR (1) | |
| <i>Pheucticus aureoventris</i> Black-backed Grosbeak | T | 1,3,5,6 | L | 1,000 | C | |
| <i>Parkerthraustes uernalis</i> Yellow-shouldered Grosbeak | T | 1,4-6 | L | 800 | R | |
| <i>Saltator rossus</i> Slate-coloured Grosbeak | T | 1,4,5 | L | 1,400 | R | |
| <i>Saltator maximus</i> Buff-throated Saltator | T | 1-6 | L | 1,400 | F | |
| <i>Saltator coerulescens</i> Greyish Saltator | O | 1,3,4 | L | 600 | C | |
| <i>Cyanocopsa cyanooides</i> Blue-black Grosbeak | T | 1-6 | L | 800 | C | |
| <i>Cyanocopsa brissonii</i> Ultramarine Grosbeak | O | 3 | 700 | 700 | VR (1) | |
| <i>Cissopis leveriana</i> Magpie Tanager | O | 1,2,4-6 | 0 | 1,300 | F | |
| <i>Chlorospingus ophthalmicus</i> Common Bush Tanager | T | 5,6 | 900 | 1,450 | C | |
| <i>Thlypopsis sordida</i> Orange-headed Tanager | O | 5 | 900 | 900 | VR (1) | |
| <i>Hemitithraupis guira</i> Guira Tanager | T | 1,5 | L | 1,000 | C | |

Appendix 2: Continued.

| Species | Evidence | Sites | Min | Max | Abundance | Breeding Information |
|---|----------|---------|-------|-------|-----------|----------------------|
| <i>Hemitraupis flavicollis</i> Yellow-backed Tanager | T | 1,4-6 | L | 1,000 | F | |
| <i>Nemosia pileata</i> Hooded Tanager | O | 1 | L | 800 | R | |
| <i>Chlorothraupis carmioli</i> Carmiol's Tanager | T | 1-6 | L | 1,200 | C | |
| <i>Lanio versicolor</i> White-winged Shrike Tanager * | T | 1,4-6 | L | 1,000 | F | |
| <i>Creurgops dentata</i> Slaty Tanager **, ** | O | 6 | 1,400 | 1,400 | VR (1) | |
| <i>Tachyphonus cristatus</i> Flame-crested Tanager | O | 5 | 600 | 600 | VR (1) | |
| <i>Tachyphonus rufiventer</i> Yellow-crested Tanager * | O | 1,4-6 | L | 1,000 | F | |
| <i>Tachyphonus luctuosus</i> White-shouldered Tanager | T | 1,3-6 | L | 1,000 | F | |
| <i>Trichothraupis melanops</i> Black-goggled Tanager | O | 5,6 | 900 | 1,200 | R | |
| <i>Habia rubica</i> Red-crowned Ant-Tanager | T | 1-6 | L | 1,000 | C | |
| <i>Piranga flava</i> Hepatic Tanager | T | 1,3,5 | 700 | 1,000 | F | NA 15-11-99 |
| <i>Piranga rubra</i> Summer Tanager | O | 4,5 | L | 1,800 | U (Mn) | |
| <i>Piranga olivacea</i> Scarlet Tanager | O | 4,5 | 700 | 1,000 | F (Mn) | |
| <i>Piranga leucoptera</i> White-winged Tanager | T | 5,6 | 700 | 1,200 | F | |
| <i>Ramphocelus carbo</i> Silver-beaked Tanager | T | 1-6 | L | 1,400 | C | |
| <i>Thraupis episcopus</i> Blue-grey Tanager | O | 1,4 | L | 900 | U | |
| <i>Thraupis sayaca</i> Sayaca Tanager | O | 1,2,5,6 | L | 1,400 | C | |
| <i>Thraupis palmarum</i> Palm Tanager | T | 1-6 | L | 1,400 | C | NB 20-11-99 |
| <i>Anisognathus flavinucha</i> Blue-winged Mountain Tanager | O | 6 | 1,300 | 1,400 | U | |
| <i>Euphonia chlorotica</i> Purple-throated Euphonia | T | 1,3,5,6 | L | 700 | C | |
| <i>Euphonia lanirostris</i> Thick-billed Euphonia | T | 1,3-6 | L | 1,200 | F | NM 20-11-99 |
| <i>Euphonia cyanocephala</i> Blue-hooded Euphonia | O | 1 | L | L | R (?Ms) | |
| <i>Euphonia chrysopasta</i> White-lored Euphonia | T | 1,4-5 | L | 1,000 | U | NM 02-08-97 |
| <i>Euphonia mesochrysa</i> Bronze-green Euphonia | T | 1,4-6 | 600 | 1,200 | C | |
| <i>Euphonia minuta</i> White-vented Euphonia | T | 1,4 | L | 1,000 | U | |
| <i>Euphonia xanthogaster</i> Orange-bellied Euphonia | T | 4-6 | L | 1,400 | C | |
| <i>Euphonia rufiventris</i> Rufous-bellied Euphonia | T | 1-6 | L | 1,000 | F | |
| <i>Chlorophonia cyanea</i> Blue-naped Euphonia | T | 1,5,6 | L | 1,400 | U | |
| <i>Chlorochrysa calliparva</i> Orange-eared Tanager | O | 6 | 1,300 | 1,300 | VR (1) | |
| <i>Tangara mexicana</i> Turquoise Tanager | T | 1,4-6 | L | 1,300 | C | NM 06-10-98 |
| <i>Tangara chilensis</i> Paradise Tanager | T | 1,3-6 | L | 1,300 | C | NM 17-11-99 |

Appendix 2: Continued.

| Species | Evidence | Sites | Min | Max | Abundance | Breeding Information |
|--|----------|-------|-------|-------|-----------|----------------------|
| <i>Tangara schrankii</i> Green-and-gold Tanager | T | 1-6 | L | 1,300 | C | F 05-02-98 |
| <i>Tangara arthus</i> Golden Tanager | O | 5,6 | 700 | 1,400 | R | |
| <i>Tangara xanthocephala</i> Saffron-crowned Tanager | O | 6 | 1,200 | 1,400 | U | |
| <i>Tangara xanthogastra</i> Yellow-bellied Tanager | T | 1,3-6 | L | 1,200 | F | |
| <i>Tangara chrysolis</i> Golden-eared Tanager | O | 6 | 1,200 | 1,400 | VR (2) | |
| <i>Tangara gyrola</i> Bay-headed Tanager | T | 1-6 | L | 1,200 | C | NM 17-11-99 |
| <i>Tangara ruficervix</i> Golden-naped Tanager | O | 6 | 1,300 | 1,300 | VR (1) | |
| <i>Tangara cyanicollis</i> Blue-necked Tanager | O | 6 | 1,300 | 1,350 | VR (2) | |
| <i>Tangara nigrocincta</i> Masked Tanager | T | 1,3-6 | L | 1,200 | C | F 16-11-99 |
| <i>Tangara nigroviridis</i> Beryl-spangled Tanager | O | 5 | 1,000 | 1,000 | VR (1) | |
| <i>Dacnis lineata</i> Black-faced Dacnis | T | 1,3-6 | L | 1,250 | F | |
| <i>Dacnis flaviventris</i> Yellow-bellied Dacnis | O | 1,4,5 | L | 800 | F | |
| <i>Dacnis cayana</i> Blue Dacnis | T | 1,3-6 | L | 1,300 | C | |
| <i>Chlorophanes spiza</i> Green Honeycreeper | T | 1,4-6 | 0 | 1,400 | F | |
| <i>Cyanerpes caeruleus</i> Purple Honeycreeper | T | 1,4-6 | 0 | 1,000 | C | |
| <i>Cyanerpes cyaneus</i> Red-legged Honeycreeper | T | 1,5 | 0 | 1,200 | F | F 07-02-98 |
| <i>Diglossa glauca</i> Deep-blue Flowerpiercer | O | 6 | 1,200 | 1,200 | VR (1) | |
| <i>Tersina viridis</i> Swallow Tanager | T | 1,3-6 | L | 1,400 | U | |
| <i>Parula pitiayumi</i> Tropical Parula | T | 1,3-6 | 0 | 1,300 | C | |
| <i>Myioborus miniatus</i> Slate-throated Redstart | T | 2,4-6 | 600 | 1,400 | C | NM 06-08-97 |
| <i>Basileuterus bivittatus</i> Two-banded Warbler | T | 4-6 | 700 | 1,400 | C | |
| <i>Basileuterus coronatus</i> Russet-crowned Warbler | T | 4 | 800 | 900 | F | |
| <i>Basileuterus culicivorus</i> Golden-crowned Warbler | T | 1-4 | L | 900 | F | F 19-10-98 |
| <i>Basileuterus tristriatus</i> Three-striped Warbler | T | 3,6 | 800 | 1,400 | F | |
| <i>Basileuterus fulvicauda</i> Buff-rumped Warbler | T | 3 | 600 | 600 | R? | |
| <i>Basileuterus rivularis</i> River Warbler | T | 1-6 | L | 900 | F? | F 20-07-97 |
| <i>Controstrum speciosum</i> Chestnut-vented Conebill | O | 1 | 450 | 450 | VR (1) | |
| <i>Controstrum albifrons</i> Capped Conebill | O | 4 | 700 | 700 | VR(1) | |
| <i>Psarocolius decumanus</i> Crested Oropendola | T | 1-6 | L | 1,300 | C | NA 20-10-98 |
| <i>Psarocolius atrovirens</i> Dusky-green Oropendola * | T | 6 | 400 | 1,400 | C | NB 04-11-98 |

References

- Arribas, M. A., Jammes, L. and Sagot, F. (1995) *Lista de las aves de Bolivia*. Santa Cruz: Armonía.
- BirdLife International (2000) *Threatened birds of the world*. Barcelona and Cambridge, U.K.: Lynx Edicions and BirdLife International.
- Haselmayer, J. and Quinn, J. S. (2000) A comparison of point counts and sound recording as bird survey methods in Amazonian southeast Peru. *Condor* 102: 887–893.
- Hennessey, A. B., Herzog, S. K., Sagot, F. (2003) *Lista Anotada de las Aves de Bolivia*. Quinta edición. Santa Cruz de la Sierra, Bolivia: Asociación Armonía/BirdLife International.
- Herzog, S. K., Soria A., R., Troncoso J., A. and Matthysen, E. (submitted MS) Composition and spatial structure of avian mixed-species flocks in a high-Andean *Polylepis* forest in Bolivia. *Ecotropica*.
- Hilty, S. L. and Brown, W. L. (1986) *A guide to the birds of Colombia*. Princeton: Princeton University Press.
- Jullien, M. and Thiollay, J. M. (1998) Multi-species territoriality and dynamic of neotropical forest understorey bird flocks. *J. Anim. Ecol.* 67: 227–252.
- Kessler, M. (2001). Diversidad y endemismo de grupos selectos de plantas en la Serranía de Pilón Lajas, depto. Beni, Bolivia. *Revista de la Sociedad Boliviana de Botánica* 3: 124–145.
- Mayer, S. (2000) *Birds of Bolivia 2.0* (CD-ROM). Westernland: Bird Songs International.
- McNair, D. B. and Escobar, J. (1993) Verified winter record of Scarlet Tanager in Charleston, SC, and a review of the winter status of Scarlet Tanager in the Southeast United States. *Chat* 57: 25–31.
- Munn, C. A. (1985) Permanent canopy and understory flocks in Amazonia: species composition and population density. *Orn. Monogr.* 36: 683–712.
- Parker, T. A., III. (1989) An avifaunal survey of the Chimanes Ecosystem Program Area of northern Bolivia. Unpublished report.
- Parker, T. A., III. (1991) On the use of tape recorders in avifaunal surveys. *Auk* 108: 443–444.
- Parker, T. A., III, and Bailey, B. (1991) *A Biological Assessment of the Alto Madidi Region and adjacent areas of Northwest Bolivia, May 18–June, 1990*. Washington, D.C.: Conservation International (RAP Working Papers 1).
- Parker, T. A., III, Remsen, J. V., Jr., and Heindel, J. A. (1980) Seven bird species new to Bolivia. *Bull. Brit. Orn. Cl.* 100: 160–162.
- Parker, T. A., III., Castillo U., A., Gell-Mann, M. and Rocha O., O. (1991) Records of new and unusual birds from northern Bolivia. *Bull. Brit. Orn. Cl.* 111: 120–138.
- Parker, T.A., III, Bates, J.M. and Cox, G. (1992) Rediscovery of the Bolivian Recurvebill with notes on other little-known species of the Bolivian Andes. *Wilson Bull.* 104: 173–178.
- Parker, T. A., Donahue, P. K. and Schulenberg, T. S. (1994) Birds of the Tambopata Reserve (Explorer's Inn Reserve). In R. B. Foster, T. A. Parker III, A. H. Gentry, L. H. Emmons, A. Chicchón, T. S. Schulenberg, L. Rodríguez, G. Lamas, H. Ortega, J. Icochea, W. Wust, M. Romo, J. A. Castillo, O. Phillips, C. Reynal, A. Kratter, P. K. Donahue and L. J. Barkley, eds. *The Tambopata Reserved Zone of southeastern Peru: a biological assessment*. Washington, D.C.: Conservation International (RAP Working Papers 6).
- Parker, T. A., III., Stotz, D. F. and Fitzpatrick, J. W. (1996) Ecological and distributional databases. In D. F. Stotz, Fitzpatrick, J.W., Parker, T. A., III, and Moskovits, D.K. *Neotropical birds: ecology and conservation*. Chicago: University of Chicago Press.
- Perry, A., Kessler, M. and Helme, N. (1997) Birds of the central Río Tuichi Valley, with emphasis on dry forest, Parque Nacional Madidi, dpto. La Paz, Bolivia. *Orn. Monogr.* 48: 557–576.

- Remsen, J. V., Jr., and Traylor, M. A., Jr. (1989) *An annotated list of the birds of Bolivia*. Vermillion, SD: Buteo Books.
- Sagot, F. (1998) Nuevas especies en el país en relación a la Lista de las Aves de Bolivia de Armonía de 1995. Pp. 25–31 in F. Sagot and J. Guerrero, eds. *Aves y conservación en Bolivia No 1 (Actas IV Enc. Bol. Cons. Aves, 25–27 Oct. 97, Tarija)*. Santa Cruz: Armonía.
- Smith, D. N. and Killeen, T. J. (1998) A comparison of the structure and composition of montane and lowland tropical forest in the Serranía Pílon Lajas, Beni, Bolivia. Pp. 681–700 in F. Dallmeier and J. A. Comiskey, eds. *Forest biodiversity in North, Central and South America and the Caribbean. Research and Monitoring*. Paris and New York: UNESCO and Parthenon Publishing Group.
- Seidel, R. (1995) Inventario de los árboles en tres parcelas de bosque primario en la Serranía de Marimonos, Alto Beni. *Ecología en Bolivia* 25: 1–35.
- Stattersfield, A. J., Crosby, M. J., Long, A. J. and Wege, D. C. (1998) *Endemic Bird Areas of the world*. Cambridge, U.K.: BirdLife International.
- Stotz, D. F., Fitzpatrick, J. W., Parker, T. A., III, and Moskovits, D.K. (1996) *Neotropical birds: ecology and conservation*. Chicago: University of Chicago Press.
- Thomas B.T. (1999) Family: Steatornis. Pp 244–251 in J. Del Hoyo, A. Elliot and J. Sargatal, eds. *The handbook of the birds of the world*, V. Barcelona: Lynx Editions

A. BENNETT HENNESSEY

Armonía/ BirdLife Int., Casilla 3566, Santa Cruz, Bolivia. E-mail: tangara@unete.com

SEBASTIAN K. HERZOG

Institut für Vogelforschung "Vogelwarte Helgoland", An der Vogelwarte 21, 26386 Wilhelmshaven, Germany.

MICHAEL KESSLER

Albrecht-von-Haller-Institut für Pflanzenwissenschaften, Abteilung Systematische Botanik, Untere Karspüle 2, 37073 Göttingen, Germany.

DANIEL ROBISON

Rurrenabaque, La Paz, Bolivia.

Received April 2002; revision accepted 9 June 2003

